

Cite this: *RSC Adv.*, 2015, 5, 78047

On the investigation of hybrid quinones: synthesis, electrochemical studies and evaluation of trypanocidal activity†

Guilherme A. M. Jardim,^a Wallace J. Reis,^a Matheus F. Ribeiro,^a Flaviano M. Ottoni,^b Ricardo J. Alves,^b Thaissa L. Silva,^c Marília O. F. Goulart,^c Antonio L. Braga,^d Rubem F. S. Menna-Barreto,^e Kelly Salomão,^e Solange L. de Castro^e and Eufrânio N. da Silva Júnior^{*a}

In our continued search for novel trypanocidal compounds, arylamine, chalcone, triazolic, triazole-carbohydrate and chalcogenium derivatives containing a naphthoquinone scaffold were prepared; in addition to electrochemical studies, these compounds were evaluated against the infective bloodstream form of *Trypanosoma cruzi*, the etiological agent of Chagas disease. Among the thirty-eight compounds herein evaluated, six were found to be more potent against trypomastigotes than the standard drug benznidazole, with IC₅₀/24 h values between 52.9 and 89.5 μM.

Received 12th August 2015
Accepted 8th September 2015

DOI: 10.1039/c5ra16213k

www.rsc.org/advances

Introduction

Chagas disease is caused by the protozoan *Trypanosoma cruzi* (*T. cruzi*) and affects approximately 5.7 million individuals in Latin America, of whom 30–40% either have or will develop cardiomyopathy, digestive megasyndromes or both.¹ The transmission of Chagas disease occurs through the faeces of sucking triatominae insects, blood transfusions, organ transplantation, oral contamination through laboratory accidents and congenital routes.² A current major concern is related to outbreaks of acute Chagas disease associated with the ingestion of contaminated food and drink.^{3,4} Another concern is the disease's emergence in non-endemic areas such as North America and Europe, a phenomenon that is likely due to the immigration of infected individuals.^{5,6} This disease is characterised by two clinical phases: a short, acute phase defined by patent parasitaemia and a long, progressive chronic phase.⁷

The life cycle of *T. cruzi* involves obligatory passage through vertebrate and invertebrate hosts in a series of different developmental stages.⁸ The trypomastigote is ingested by the insect and differentiates into epimastigote form, which differentiates into the metacyclic trypomastigote form. Following the invasion of vertebrate host cells, this form differentiates into the

amastigote, which after several reproductive cycles, transforms into the trypomastigote form, which is responsible for dissemination of the infection.

The available chemotherapy for this illness includes two nitroheterocyclic agents, nifurtimox and benznidazole (Bnz). These two agents are effective against acute infections but show poor activity, severe collateral effects and limited efficacy against different parasitic isolates in the late chronic phase. Thus, there is an urgent need to identify better drugs to treat chagasic patients.^{9,10}

Among several naturally occurring compounds, the naphthoquinones have emerged with a broad distribution in the plant kingdom; naphthoquinones are involved in oxidative processes due to their structural properties and their capacity to generate reactive oxygen species.^{11,12} Quinoidal compounds can be found in various plant families or as synthetic substances^{13,14} and are the focus of research aimed at the identification of derivatives with efficacy and selectivity toward *T. cruzi*.¹⁵

Lapachol (**1**) is an important natural naphthoquinone; our group has focused on synthesising and evaluating the trypanocidal activity of lapachones substituted with 1,2,3-triazole moieties.¹⁵ The broad and potent activity of triazole and their derivatives has established them as pharmacologically significant scaffolds.^{16,17} The catalysis by Cu(I) of 1,3-dipolar cycloadditions between azides and terminal alkynes¹⁸ led to enhanced reactivity and regioselectivity, resulting in the formation of 1,4-disubstituted 1,2,3-triazoles.¹⁹ This conversion has been used increasingly in drug discovery due to its reliability, specificity, and biocompatibility; also, triazoles can act as both a bioisostere and a linker that readily associates with biological targets through hydrogen-bonding and dipole interactions.²⁰

^aInstitute of Exact Sciences, Department of Chemistry, UFMG, Belo Horizonte, MG, 31270-901, Brazil. E-mail: eufranio@ufmg.br

^bSchool of Pharmacy, UFMG, Belo Horizonte, MG, 31270-901, Brazil

^cInstitute of Chemistry and Biotechnology, UFAL, Maceió, AL, 57072-970, Brazil

^dDepartment of Chemistry, UFSC, Florianópolis, SC, 88040-900, Brazil


^eLaboratory of Cellular Biology, IOC, FIOCRUZ, Rio de Janeiro, RJ, 21045-900, Brazil

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c5ra16213k


Over the years, our group has investigated the synthesis, selectivity, and mechanisms of action of quinoidal compounds in relation to *T. cruzi*.²¹ The synthetic strategies involved A-ring, C-ring and redox centre modifications of lapachones.^{15,21,22} For instance, naphthoimidazoles were prepared by redox centre modification of β -lapachone; among the derivatives assayed, three compounds were active against bloodstream trypomastigotes of *T. cruzi* (Scheme 1a).^{23,24} 3-Arylamino nor- β -lapachones represent an important example of C-ring-modified lapachones. The compounds were effective against trypomastigote forms,

with nor- β -lapachone dibromo-substituted arylamine exhibiting approximately four times more activity than Bnz, the standard drug (Scheme 1a).²⁵ Nor- β -lapachone-based 1,2,3-triazole (which is two times more active than Bnz) was also effective against the proliferative forms of *T. cruzi* (Scheme 1a).^{26,27} Silva and co-workers²⁸ described the synthesis of iodinated and methylated naphthoquinones obtained from C-allyl lawsone (Scheme 1a) that were active against epimastigotes and trypomastigotes of *T. cruzi*, with the main target being the mitochondrion.²⁹


a) Trypanocidal compounds obtained by redox centre or C-ring modifications



b) Examples of carbohydrate and chalcone hybrids with trypanocidal activity



c) Quinones evaluated in this work against *T. cruzi*.



Scheme 1 Overview. * $\text{IC}_{50}/24 \text{ h}$ values for the lytic activity on bloodstream trypomastigotes.

The triazole moiety is also used for the synthesis of glycosylated structures.³⁰ For example, a series of sugar triazoles has been synthesized, and 1,2,3-triazole-linked sialic acid-6-*O*-galactose was active *in vitro* against *T. cruzi*.³¹ Recently, Campo and co-workers described 1,4-disubstituted 1,2,3-triazole-sialic acid derivatives containing galactose, glucose or gulose as potential inhibitors of *T. cruzi* trans-sialidase (Scheme 1b).³²

Chalcones can be readily synthesized by the Claisen-Schmidt condensation method,³³ and they represent a key moiety in medicinal chemistry, displaying a wide range of pharmacological activities.³⁴ Hybrids of chalcone with quinoline,³⁵ paullone³⁶ and benzoxaborole³⁷ have shown significant activity against pathogenic trypanosomatids and low cytotoxicity to mammalian cells (Scheme 1b).

Finally, several reports have described chalcogen-containing quinones with antibacterial, antifungal³⁸ and antitumor activities,³⁹ and the last two decades have witnessed a growing interest in selenium-containing compounds with potential 'redox modulating' properties. Certain redox catalysts containing quinone- and chalcogen-bearing building blocks have shown considerable importance, with activity against cancer cells and other biological targets.⁴⁰

In this context, we herein describe the synthesis and trypanocidal activity of (i) α -lapachone arylamines: as discussed above, the presence of an arylamino group in nor- β -lapachone improves the trypanocidal activity, and the biological activity of 4-arylamino α -lapachone remains unexplored; (ii) nor- β -lapachone-based chalcones: we aim to obtain active compounds, due to the well-known bioactivity of each individual moiety; (iii) quinone-based 1,2,3-triazoles from C-allyl lawsone: the insertion of 1,2,3-triazole could enhance the trypanocidal activity of the derivatives previously shown to be less active than Bnz, as displayed in Scheme 1a;¹⁵ (iv) naphthoquinone-based 1,2,3-triazole


linked to monosaccharides: sugar residues are able to modulate physicochemical parameters, such as solubility and interactions with cell surface components, which are both important parameters enabling a drug to act on intracellular parasites; and (v) selenium- and sulphur-containing lapachones: for the first time, this class of compounds is evaluated against *T. cruzi* and subjected to concise electrochemical studies (Scheme 1c).

Results and discussion


Chemistry

To obtain α -lapachone arylamines, α -lapachone (**2**) was prepared in an initial step from lapachol (**1**) in acidic conditions.⁴¹ Then, **2** was reacted with NBS in CCl₄ using benzoyl peroxide as an initiator to afford 4-bromo- α -lapachone; a nucleophilic substitution reaction with substituted anilines gave the compounds **3–14** (Scheme 2). We recently reported compounds **3–7**, **9–10** and **14**,⁴² and compounds **8**, **11–13** are herein described for the first time. In order to achieve nor- β -lapachone-based chalcones, amino-chalcones were synthesized *via* a Claisen condensation reaction with 4'-aminoacetophenone and different aldehydes in the presence of sodium hydroxide in ethanol.⁴³ Then, nor-lapachol (**15**)⁴⁴ was treated with excess bromine to obtain 3-bromo-nor- β -lapachone (**16**) in quantitative yield.⁴¹ Through a nucleophilic substitution reaction with amino-chalcones and **16**, compounds **17–21** were prepared as previously published (Scheme 2).⁴⁵

To obtain the third class of substances, C-allyl-lawsone (**22**) was prepared in an initial step from the reaction of lawsone with allyl bromide.⁴⁶ In a second step, **22** was treated with excess iodine,²⁸ forming the *para* and *ortho* regioisomers. The iodinated naphthoquinone **23** (*ortho*-derivative) was obtained in moderate yield as a crystalline red solid after separation by column chromatography. Compound **24** was synthesized in




Scheme 2 Synthesis of α -lapachone arylamines and nor- β -lapachone-based chalcones.



Scheme 3 Synthesis of quinone-based 1,2,3-triazoles 26–30 and linkage to monosaccharides to give 30 and 31. Conditions (i): $CuSO_4 \cdot 5H_2O$, sodium ascorbate, CH_2Cl_2/H_2O .

good yield *via* nucleophilic substitution of 23 with sodium azide in dimethylformamide (DMF), as previously described.⁴⁷ The 1,2,3-triazole derivatives were obtained through a click reaction

of 24 with $CuSO_4 \cdot 5H_2O$ /sodium ascorbate and the corresponding alkynes in a mixture of CH_2Cl_2/H_2O , giving 25–29. Alkyne derivatives of monosaccharides were also used in the



Scheme 4 Synthesis of nor- β -lapachone-based 1,2,3-triazole-carbohydrates 33–35.

click reaction, providing **30–31** (Scheme 3). These alkyne-carbohydrate derivatives were obtained using two different methods of acetylation.^{48,49} For the glucose and galactose derivatives, D-glucose and D-galactose were subjected to peracetylation with NaOAc in acetic anhydride, and the compounds were prepared in moderate yields. For the mannose derivative, catalytic amounts of I₂ were used instead of NaOAc. The product was reacted with propargyl alcohol and catalytic amounts of BF₃·Et₂O in CH₂Cl₂ to achieve the alkyne derivative.⁵⁰

To prepare nor-β-lapachone-based monosaccharide-linked 1,2,3-triazoles, bromo derivative **16** was subjected to a nucleophilic substitution reaction with sodium azide in dichloromethane to obtain **32** in quantitative yield, as previously reported.²⁶ From the reaction of **32** with CuSO₄·5H₂O/sodium ascorbate and the alkyne derivatives of glucose, mannose and galactose in a mixture of CH₂Cl₂/H₂O, compounds **33–35** were obtained in good yields (Scheme 4).

Finally, as recently described,⁵¹ the chalcogenylation of lapachol (**1**) was induced under microwave irradiation and using molecular iodine as a catalyst, DMSO as a stoichiometric oxidant and different chalcogenium compounds, affording the selenium- and sulphur-containing β-lapachone derivatives (Scheme 5).

The structures of the novel compounds **8**, **11–13**, **25–31** and **33–35** were determined by ¹H and ¹³C NMR. Electrospray ionization mass spectra were also obtained. The ¹H and ¹³C NMR spectra for all unpublished compounds are included in the ESI.†

Biological and electrochemical analysis


The compounds **3–14**, **17–21**, **25–31** and **33–46** were evaluated against trypomastigote forms of *T. cruzi*. The more active

compounds against *T. cruzi* were **14**, **26–29** and **41**, which exhibited IC₅₀/24 h values in the range of 52.9 to 89.5 μM, in comparison with the IC₅₀/24 h value of 103.6 μM for the reference drug, Bnz. Thus, in general, these compounds are approximately two times more potent than Bnz (Table 1).

Table 1 Activity of compounds against the trypomastigote form of *T. cruzi*

Compd	IC ₅₀ /24 h ^a (μM)	Compd	IC ₅₀ /24 h ^a (μM)
1	410.8 ± 53.2 ^b	26	89.5 ± 2.8
2	>4000 ^b	27	52.9 ± 2.1
3	>4000	28	55.3 ± 5.2
4	911.0 ± 42.2	29	86.7 ± 5.4
5	374.5 ± 179.5	30	300.3 ± 21.1
6	>4000	31	174.9 ± 10.6
7	>4000	32	50.2 ± 3.8 ^d
8	1471.3 ± 171.6	33	472.2 ± 16.5
9	>4000	34	208.9 ± 7.3
10	1093.8 ± 133.1	35	202.3 ± 6.0
11	>4000	36	102.9 ± 0.0
12	440.6 ± 73.0	37	1677.8 ± 113.3
13	>4000	38	207.6 ± 33.52
14	89.3 ± 6.0	39	360.5 ± 34.6
15	1280.6 ± 167.2 ^b	40	477.1 ± 14.83
17	>4000	41	54.9 ± 3.19
18	>4000	42	251.9 ± 52.51
19	787.4 ± 40.6	43	>4000
20	230.2 ± 71.6	44	281.8 ± 3.45
21	>4000	45	109.2 ± 33.2
22	352.0 ± 29.0 ^c	46	>4000
23	398.0 ± 56.0 ^c	Benznidazole ^d	103.6 ± 0.6
25	213.5 ± 8.4	Crystal violet ^b	536.0 ± 3.0

^a Mean ± SD of at least three independent experiments. ^b Ref. 23c. ^c Ref. 28. ^d Ref. 26.



Scheme 5 Chalcogenium-containing β-lapachone derivatives **36–46** evaluated against *T. cruzi*.

We recently described the arylamino-substituted α -lapachone compounds **3–14** as having moderate antitumor activity.⁴² Herein, we prepared a series of arylamino-substituted α -lapachone compounds possessing electron-withdrawing and electron-donating groups. No clear structure–activity relationship was observed, and all the compounds were either inactive or presented low activity, with the exception of compound **14** ($IC_{50}/24$ h of 89.3 μ M).

As discussed before, chalcones represent an important class of compounds with trypanocidal activity.^{33–37} Based on this previously determined biological activity, we prepared and evaluated the nor- β -lapachone-based chalcone hybrids **17–21**. These compounds were previously shown to possess potent antitumor activity.⁴⁵ Unfortunately, the compounds failed to kill *T. cruzi*, and the derivatives were considered inactive; the exception was compound **20** ($IC_{50}/24$ h of 230.2 μ M), which presented moderate activity against trypomastigote forms of the parasite.

The third class of compounds, lapachone derivatives obtained from C-allyl lawsone (**22**), are promising trypanocidal agents. We recently used molecular hybridization of naphthoquinoidal compounds and 1,2,3-triazoles to design trypanocidal derivatives.^{25,26,52} We discovered, for instance, four classes of naphthoquinone-based 1,2,3-triazoles, which are exemplified by compounds **47–52** (Fig. 1), which have the pronounced ability to inhibit the infective bloodstream form of *T. cruzi* and are more active than Bnz. We described some insights about the mechanism of action of **48**. This substance induced reservosome disruption, flagellar blebbing and Golgi disorganization in the parasites. According to our studies, autophagy could be a part of the mechanism of action for this derivative.⁵³

In our search for novel quinone-based 1,2,3-triazoles, we herein describe for the first time a series of triazolic derivatives prepared from C-allyl lawsone (**22**) (Scheme 3). Compounds **26**, **27**, **28** and **29**, with $IC_{50}/24$ h values of 89.5, 52.9, 55.3 and 86.7 μ M, respectively, presented important trypanocidal activity and

will be considered for further studies. These derivatives were obtained in only three steps from C-allyl lawsone (**22**), and **27** and **28** were almost twice as active as Bnz. Additionally, we prepared lapachone coupled with carbohydrate derivatives **30**, **31** and **33–35** to evaluate the ability of these compounds to generate damage to the parasite. We observed that these new compounds presented moderate activity against *T. cruzi*, but the compounds were less active than Bnz. In general terms, the strategy was efficient, but further structural modifications will be required to improve the biological potential of these compounds.

We recently described the last class of compounds as possessing potential antitumor activity.⁵¹ Selenium- and sulphur-containing β -lapachone derivatives **33–46** were active against *T. cruzi* with generally moderate activity; however, compounds **41** ($IC_{50}/24$ h = 54.9 μ M), **36** ($IC_{50}/24$ h = 102.9 μ M) and **45** ($IC_{50}/24$ h = 109.2 μ M) exhibited IC_{50} values similar to that of Bnz, and **41** was 2-fold more potent than Bnz. Herein, we describe detailed studies of the electrochemical aspects of these compounds, as only a few examples of chalcogenium-containing lapachones have been reported, along with evaluation of anticancer activity. To the best of our knowledge, this is the first joint report of electrochemistry and trypanocidal activity results, related to this class of compounds.

In addition, we also evaluated the cytotoxicity of the most active compounds against peritoneal macrophages. The results showed that the active substances were more toxic to the parasites than to the cell lines tested. Finally, our results demonstrated that compounds **14** and **26–29** are the most active against this trypanosomatid and may represent promising compounds for treatment of Chagas disease.

Molecular electrochemistry, through non-destructive techniques such as cyclic voltammetry in aprotic and protic media, has proved to be very useful for characterizing electron transfers and deciphering chemical reaction mechanisms that are associated with electron transfer, with some correlation with redox-based biological activity.^{12,54} In this context, our group has described the synthesis and electrochemical studies of several quinones in an attempt to correlate them with trypanocidal activity,⁵⁵ as well as the electrochemical aspects of arylamino-substituted α -lapachones and nor- β -lapachones.^{42,52,56}

Among all the investigated multifunctional compounds herein described, only a few examples of electrochemical studies on chalcogen-containing quinones have been reported in the literature.⁴⁰ In order to corroborate the biological aspects of selenium- and sulphur-containing β -lapachones and establish more insight using electrochemical parameters, we performed experiments with the compounds **36–39**, **40**, **41** and **45**. Using cyclic voltammetric studies in aprotic (DMF/TBAPF₆) and protic (ethanolic phosphate buffered medium at pH 7.4) media, the two redox centres (quinoidal and organochalcogen moieties) were investigated.⁵⁷ The electrochemical properties of organic selenium compounds^{58,59} qualitatively resemble those of the more abundant sulphur compounds, although they are much less explored. The number of redox potentials of organoselenium organic compounds is limited. The products of electrochemical oxidation are remarkably dependent upon the




Fig. 1 Trypanocidal compounds designed by molecular hybridization of quinones and 1,2,3-triazoles. * $IC_{50}/24$ h values for the lytic activity on bloodstream trypomastigotes.

structure of the organic moiety, the water content of the solvent, and the supporting electrolyte.⁵⁸

In the present case, cyclic voltammogram (CV) was used to investigate the electrochemical behaviour of the seven selenoquinones **36–39**, **40**, **41**, and **45**, which were recently reported as antitumor compounds.⁵¹ The obtained profiles confirmed the presence of two individual redox centres.

In typical measurements, CVs were recorded in aprotic medium (DMF + TBAPF₆, 0.1 mol L⁻¹) to resemble the membrane environment at a scan rate of 100 mV s⁻¹. This enabled the determination of the electrochemical reduction (from 0.5 up to -3.0 V) and oxidation (from -0.5 up to 2.0 V) behaviours of each compound. Cathodic and anodic peak potentials for each compound are listed in Table 2. All the compounds showed electrochemical activity in the cathodic and anodic arms of the CVs, as represented by compound **38** (Fig. 2).

The voltammograms for compounds **36–39**, **40**, **41** and **45** displayed similar profiles in the cathodic and anodic regions. Fig. 3 shows the electrochemical behaviour of compound **40** as a representative of the series.

As expected, the overall CV profiles for the quinonoid compounds are similar to those reported for other quinones.^{42,53,60} One couple of cathodic and anodic peaks is represented by a diffusional ($E_{\text{pic}} \propto \nu^{1/2}$) quasi-reversible nature (Ic/Ia). This first pair is related to the anion-radical (semi-quinone) formation, and the second pair of peaks is broader and ill-defined (IIc/IIa), as observed before,⁶⁰ due to a possible disproportionation of *ortho*-quinones (Fig. 3a and b). In the anodic region, a sharp, irreversible peak (IVa) appears, with at least a duplicated current intensity. In the reversed cycle, a reduction wave (IVc) is observed close to 0 V; in the successive scan, it is observed that the reduced product formed can be re-oxidized (IVa') (Fig. 2c). In the case of compound **38**, anodic peak IVa is followed by a more positive one (Va) due to the presence of an electron-donating group. Electrochemical data are displayed in Table 2.

The values of E_{pic} varied from -0.644 V up to -0.674 V (Table 2). As the selenium-phenyl moiety is coupled to the reducible quinone moiety without conjugation, there is only a slight change in the reduction potentials, with compound **40** (*p*-Br) being the most easily reduced; the aliphatic selenium derivative is reduced at more negative potentials. The majority of




Fig. 2 Cyclic voltammograms (first cycle) of compound **38**, in DMF + TBAPF₆ (0.1 mol L⁻¹). Glassy carbon electrode, $\nu = 100$ mV s⁻¹, ($c = 1$ mmol L⁻¹). Black: potential range from +0.5 V to -3.0 V, cathodic direction; red: potential range from -0.5 V to +2.0 V, anodic direction.

selenium-containing β -lapachone derivatives are voltammetrically interrogated in DMF and displayed only one irreversible oxidation peak between +1.126 V and +1.252 V vs. Ag|AgCl, Cl⁻ (Table 2).


The first monoelectronic transfer leads to the formation of the corresponding Se radical-cation. Three reaction paths are generally available to alkylphenyl chalcogenide radical-cations: deprotonation; nucleophilic attack at the chalcogen atom; and C-chalcogen bond cleavage. The relative weight of each of these reaction pathways is governed by the nature of the chalcogen and of the alkyl group linked to it, as well as by the nature of the electrolysis medium.⁶¹

The electrochemical oxidation is shown to follow an overall EC-type mechanism, in which the electro-generated selenium-based cation radical undergoes an irreversible carbon-chalcogen bond rupture,⁵⁸ similar to the pathway followed by glycosylchalcogenides.⁵⁹ This cleavage produces the corresponding unsaturated quinone, which undergoes reduction in the reverse cycle (wave IVc), together with the intermediate organic selenium radical. The organic selenium radical reacts further to form the selenoxide specie, which can follow several pathways depending on the amount of water in the supporting


Table 2 Main electrochemical parameters for the analysed compounds **36–39**, **40**, **41**, and **45** in DMF + TBAPF₆ (0.1 mol L⁻¹). Glassy carbon electrode, $\nu = 100$ mV s⁻¹ ($c = 1$ mmol L⁻¹)^a

Compd	$-E_{\text{pic}}$ (V)	$-E_{\text{pic}}$ (V)	$-E_{\text{pic}}$ (V)	$-E_{\text{pic}}$ (V)	$-E_{\text{pic}}$ (V)	$-E_{\text{pic}}$ (V)	E_{pIVa} (V)	E_{pVa} (V)	E_{pIVa} (V)
36	0.655	1.134	1.340	1.269	1.063	0.548	0.301	1.126	—
37	0.655	1.061	—	0.923	0.566	0.211	0.371	1.131	—
38	0.668	1.140	—	0.945	0.580	—	0.328	1.252	1.534
39	0.658	1.140	—	1.028	0.554	0.168	0.301	1.163	—
40	0.644	1.098	—	0.913	0.542	0.132	0.337	1.198	—
41	0.662	1.123	—	0.939	0.566	0.214	0.386	1.233	—
45	0.674	1.154	—	0.937	0.584	0.247	0.309	1.137	—


^a Nd. not determined.



(a)



(b)



(c)

Fig. 3 Cyclic voltammograms of compound **40** in DMF + TBAPF₆ (0.1 mol L⁻¹). Glassy carbon electrode, $\nu = 100 \text{ mV s}^{-1}$ ($c = 1 \text{ mmol L}^{-1}$). (a) Potential range from +0.5 V to -2.0 V, 2 successive cycles. (b) Potential range from +0.5 V to -2.0 V, several values for potential inversion. (c) Potential range from -0.5 V to +1.5 V, 2 successive cycles.




Fig. 4 Cyclic voltammograms of compound **37**, in phosphate buffer (pH 7.4) + 20% ethanol. Glassy carbon electrode, $\nu = 100 \text{ mV s}^{-1}$, ($c = 0.1 \text{ mmol L}^{-1}$).

electrolyte.⁵⁷ The resulting cathodic wave IVc is a composed one, possibly related to more than one reduction pathway. The mechanism will be further explored and presented elsewhere.

The electrochemical oxidation profile, based on the values of E_{pVa} (Table 2, column 9), is related to effects of the substituents and the order of oxidation facility was found to be: **36** ~ **37** ~ **45** > **39** > **40** > **41** > **38**.

We also studied the behaviour of the compounds in protic medium. The two redox centres in compounds **36–39**, **40**, **41** and **45** were then analysed by cyclic voltammetry in protic media. The latter assessment provides insight into the oxidation and reduction processes in which compounds might participate both *in vitro* and *in vivo*. This medium resembles the hydrophilic regions of the biological matrixes.^{12,54}

The CV in protic medium is represented by a reversible peak system (Ic/Ia) in the cathodic region, typical of quinones,⁶² and one irreversible peak (IIa) with at least twice the current intensity in the anodic region. Fig. 4 represents a general electrochemical profile.

The present compounds exhibited two distinct redox behaviours that were indicative of one quinone and one chalcogen centre (Table 3). At pH 7.4, the quinone redox couple (E_{pIa} and E_{pIc}) appeared in the range of -0.260 V up to -0.288 V vs. Ag|AgCl, with quasi-reversible characteristics. Regarding the oxidation of chalcogen centres, CVs displayed irreversible oxidation peaks (E_{pIIa}) between +0.707 V (compound **36**) and

Table 3 Main electrochemical parameters for the analysed compounds in phosphate buffer (pH 7.4) + 20% ethanol. Glassy carbon electrode, $\nu = 100 \text{ mV s}^{-1}$ ($c = 0.1 \text{ mmol L}^{-1}$)

Compd	$-E_{pc_1}$ (V)	$-E_{pc_2}$ (V)	$-E_{pa_1}$ (V)	E_{pa_2} (V)	Others (V)
36	0.260	0.410	0.221	0.707	0.082
37	0.288	—	0.215	1.362	—
39	0.284	—	0.189	1.460	0.278
40	0.273	—	0.209	1.336	0.171
41	0.263	—	0.205	1.445	—
45	0.269	—	0.223	1.316	—

+1.460 V (compound **39**). The oxidation peaks are probably due to the formation of the intermediate radical species, rapidly followed by reaction with water to form the chalcogen oxide.³⁹

Overall, cyclic voltammetry confirmed the presence of two individual redox centres. While the centres influenced each other, *e.g.*, *via* electron withdrawing effects, there was little enhancement or synergy between them. This was not surprising, as there was a substantial difference between the organochalcogen and quinone redox potentials and no conjugation between them.

Conclusions

We described thirty-eight compounds that were evaluated for their effects on the infective bloodstream form of *T. cruzi*, the etiological agent of Chagas disease. Six compounds were discovered to have potent trypanocidal activity. The quinone-based 1,2,3-triazoles showed the best results in the series, with 4 representatives (**26–29**) being more active than Bnz. The strategy of preparing a wide range of multifunctional catalysts for generating oxidative stress, with two or even more redox centres combined in chemically simple molecules, was shown to be effective. Based on current assessments, structural aspects and electrochemical parameters do not show correlation with trypanocidal activity. This work represents an important contribution to the discovery of new drugs to combat *T. cruzi* and therefore provides the impetus for follow-up investigations involving synthetic chemistry, electrochemistry and cell biology.

Experimental section

General

Melting points were obtained on a Thomas Hoover apparatus and are uncorrected. Analytical grade solvents were used. Column chromatography was performed on silica gel (Silia-Flash G60 UltraPure – 60–200 μm , 60 \AA). Infrared spectra were recorded on an FTIR Spectrometer IR Prestige-21 – Shimadzu. ¹H- and ¹³C-NMR were recorded at room temperature using a Bruker AVANCE DRX200 and DRX 400 in the solvents indicated, with TMS as an internal reference. Chemical shifts (δ) are given in ppm and coupling constants (J) in Hertz. High-resolution mass spectra (electrospray ionization) were obtained using a Shimadzu LCMS-IT-TOF. Lapachol (**1**) was extracted from the heartwood of *Tabebuia* sp. (Tecoma) and purified by a series of recrystallizations. Nor-lapachol (**15**) was obtained as a crystalline orange solid by Hooker oxidation⁴⁴ of quinone **1**. Compounds **17–21** and **36–46** were prepared as recently reported by our research group.^{45,51}

Trypanocidal assay

Stock solutions of the compounds were prepared in dimethyl sulfoxide (DMSO, with the final concentration of the latter in the experiments never exceeding 0.1%). Preliminary experiments showed that concentrations of up to 0.5% DMSO have no deleterious effect on the parasites. Bloodstream

trypomastigotes of the Y strain were obtained at the peak of parasitaemia from infected albino mice, isolated by differential centrifugation and resuspended in Dulbecco's modified Eagle medium (DME) to a parasite concentration of 10^7 cells per mL in the presence of 10% of mouse blood. This suspension (100 μL was added in the same volume of each compound previously prepared at twice the desired final concentrations). Cell counts were performed in a Neubauer chamber, and the trypanocidal activity was expressed as IC_{50} , corresponding to the concentration that leads to lysis of 50% of the parasites.

Toxicity to macrophages

The cytotoxicity assays were performed using primary cultures of peritoneal macrophages obtained from Swiss mice.⁶³ For the experiments, 6×10^4 cells in 100 mL of RPMI-1640 medium (pH 7.2 plus 10% foetal bovine serum and 2 mM glutamine) were added to each well of a 96-well microtiter plate and incubated for 24 h at 37 °C. The treatment of the cultures was performed in fresh supplemented medium without phenol red (200 μL per well for 24 h at 37 °C). After this period, 110 μL of the medium was discarded and 10 μL of PrestoBlue (Invitrogen) was added to complete the final volume of 100 μL . Then, the plate was incubated for 5 h, and the measurement was performed at 570 and 600 nm, as recommended by the manufacturer. The results are expressed as the difference in the percentage of reduction between the treated and untreated cells as the LC_{50} value, corresponding to the concentration that leads to damage of 50% of the mammalian cells.

Statistical analysis

The comparison between the IC_{50} values for *T. cruzi* was performed using ANOVA followed by the Student–Newman–Keuls and Mann–Whitney tests ($p < 0.05$).

Electrochemical studies

Cyclic voltammetry (CV) experiments were performed with a conventional three electrode cell in an Autolab PGSTAT-30 potentiostat (Echo Chemie, Utrecht, the Netherlands) coupled to a PC microcomputer, using GPES 4.9 software. The working electrode was a glassy carbon electrode (GCE) BAS ($d = 3$ mm), the counter electrode was a Pt wire, and the reference electrode was an Ag|AgCl, Cl^- (sat.). All electrodes were contained in a one-compartment electrochemical cell with a volumetric capacity of 10 mL. The GC electrode was cleaned up by polishing with alumina on a polishing felt (BAS polishing kit). The solvent used was *N,N*-dimethylformamide extra dry 99.8%, acquired from Acrós Organics. In CV experiments, the scan rate varied from 10 to 500 mV s^{-1} . Electrochemical reduction was performed in aprotic media (DMF + TBAPF₆ 0.1 mol L^{-1}) at room temperature (25 ± 2 °C). Each compound (1×10^{-3} mol L^{-1}) was added to the supporting electrolyte, and the solution was deoxygenated with argon before the CV measurements. Different potential ranges were used from the cathodic to anodic scan. The most representative range was from 0 to -3.0 V vs. Ag|AgCl, Cl^- (sat.) and was chosen for the investigation

and representation of all the compounds. Anodic scans until +1.0 V were performed.

General procedure for the synthesis of compounds 3–14

To a mixture of 242 mg (1.0 mmol) of **2** and 267 mg (1.5 mmol) of *N*-bromosuccinimide in 30 mL of CCl₄, 2 mg of benzoyl peroxide was added. The mixture was heated (80 °C) for five hours. The CCl₄ was then removed by reduce pressure and 25 mL of dichloromethane and the corresponding aniline was added. The resulting mixture was stirred overnight at room temperature. The crude material was then submitted to column chromatography utilizing a mixture of ethyl acetate/hexane as an eluent.

4-((4-Fluorophenyl)amino)-2,2-dimethyl-3,4-dihydro-2H-benzo[*g*]chromene-5,10-dione (8). Compound **8** was obtained as an orange solid (210 mg, 0.6 mmol, 60% yield); mp 173–175 °C. ¹H NMR (200 MHz, CDCl₃) δ: 8.08–8.01 (m, 2H), 7.79–7.62 (m, 2H), 7.02–6.93 (m, 2H), 6.76–6.69 (m, 2H), 4.68–4.58 (m, 2H), 2.24 (dd, 1H, *J* = 3.7 and 14.3 Hz), 2.01 (dd, 1H, *J* = 5.3 and 14.3 Hz), 1.56 (s, 3H), 1.51 (s, 3H). ¹³C NMR (50 MHz, CDCl₃) δ: 184.3, 180.2, 155.8, 143.1, 134.7, 133.6, 132.4, 131.1, 128.9, 127.4, 126.7 (d, *J* = 11.2 Hz), 119.1, 116.4, 116.1 (d, *J* = 10.2 Hz), 116.0, 79.6, 65.7, 45.1, 37.8, 28.8, 26.8. ESI-MS: (*m/z*) [C₂₁H₁₈FNO₃]⁺: 351.1270. Cald. for [C₂₁H₁₈FNO₃]⁺: 351.1271.

4-((3-Iodophenyl)amino)-2,2-dimethyl-3,4-dihydro-2H-benzo[*g*]chromene-5,10-dione (11). Compound **11** was obtained as a brown solid (321 mg, 0.7 mmol, 70% yield); mp 190–192 °C. ¹H NMR (200 MHz, CDCl₃) δ: 7.97–7.93 (m, 2H), 7.67–7.63 (m, 2H), 7.04–6.96 (m, 1H), 6.88 (t, 1H, *J* = 7.9 Hz), 6.65 (d, 1H, *J* = 7.9 Hz), 4.68 (s, 1H), 4.61 (s, 1H), 2.25 (dd, 1H, *J* = 3.7 and 14.3 Hz), 1.96 (dd, 1H, *J* = 5.3 and 14.3 Hz), 1.53 (s, 6H). ¹³C NMR (50 MHz, CDCl₃) δ: 183.4, 179.7, 155.3, 148.0, 134.3, 133.2, 131.8, 130.7, 128.4, 126.9, 126.3, 126.0, 121.9, 118.3, 112.8, 95.3, 79.1, 42.9, 37.1, 29.0, 26.0. ESI-MS: (*m/z*) [C₂₁H₁₈INO₃Na]⁺: 482.0433. Cald. for [C₂₁H₁₈INO₃Na]⁺: 482.0229.

4-((3-Chlorophenyl)amino)-2,2-dimethyl-3,4-dihydro-2H-benzo[*g*]chromene-5,10-dione (12). Compound **12** was obtained as a yellow solid (220 mg, 0.6 mmol, 60% yield); mp 177–179 °C. ¹H NMR (200 MHz, CDCl₃) δ: 8.06–7.97 (m, 2H), 7.72–7.63 (m, 2H), 7.11–7.07 (m, 1H), 6.77–6.63 (m, 2H), 6.58–6.56 (m, 1H), 4.65 (s, 1H), 2.26 (dd, 1H, *J* = 2.7 and 14.4 Hz), 1.99 (dd, 1H, *J* = 5.2 and 14.4 Hz), 1.54 (s, 3H), 1.53 (s, 3H). ¹³C NMR (50 MHz, CDCl₃) δ: 183.4, 179.7, 155.3, 148.1, 135.0, 134.3, 133.2, 132.0, 130.8, 130.2, 126.3, 126.1, 118.5, 117.9, 113.1, 111.8, 79.0, 43.11, 37.4, 28.9, 26.1. ESI-MS: (*m/z*) [C₂₁H₁₈ClNO₃]⁺: 367.0962. Cald. for [C₂₁H₁₈ClNO₃]⁺: 367.0975.

4-((3-Methoxyphenyl)amino)-2,2-dimethyl-3,4-dihydro-2H-benzo[*g*]chromene-5,10-dione (13). Compound **13** was obtained as a brown solid (254 mg, 0.7 mmol, 70% yield); mp 88–90 °C. ¹H NMR (200 MHz, CDCl₃) δ: 7.92–7.91 (m, 2H), 7.70–7.53 (m, 2H), 7.06 (t, 1H, *J* = 7.9 Hz), 6.42–6.18 (m, 3H), 4.63 (s, 1H), 3.73 (s, 3H), 2.28 (dd, 1H, *J* = 2.7 and 14.4 Hz), 1.92 (dd, 1H, *J* = 5.2 and 14.4 Hz), 1.53 (s, 3H), 1.49 (s, 3H). ¹³C NMR (50 MHz, CDCl₃) δ: 183.3, 179.7, 160.7, 155.1, 148.2, 134.2, 133.1, 131.9, 130.6, 130.0, 126.2, 126.0, 118.8, 106.6, 103.0, 99.6, 79.0, 55.0,

43.0, 37.2, 28.9, 26.1. ESI-MS: (*m/z*) [C₂₂H₂₁NO₄H]⁺: 364.143. Cald. for [C₂₂H₂₁NO₄H]⁺: 364.1549.

General procedure for the synthesis of compounds 25–29

To a mixture of 255 mg (1.0 mmol) of **24**, CuSO₄·5H₂O (12.48 mg, 5 mol%) and sodium ascorbate (19.81 mg, 5 mol%) in 8 mL CH₂Cl₂/H₂O (1 : 1 v/v), the corresponding alkyne (1.1 equivalents) was added. The mixture was stirred overnight at room temperature. The organic phase was extracted with dichloromethane, dried with NaSO₄ and concentrated under reduced pressure. The residue obtained was purified by column chromatography on silica gel using as eluent a gradient mixture of hexane/ethyl acetate with increasing polarity.

2-((4-Cyclohexyl-1H-1,2,3-triazol-1-yl)methyl)-2,3-dihydronaphtho[1,2-*b*]furan-4,5-dione (25). Using ethynylcyclohexane (118 mg, 1.1 mmol), **25** was obtained as a red solid (298 mg, 0.82 mmol, 82% yield); mp 163–164 °C. ¹H NMR (400 MHz, CDCl₃) δ: 8.05 (d, 1H, *J* = 7.4 Hz), 7.70–7.66 (m, 1H), 7.62–7.58 (m, 2H), 7.54 (s, 1H), 6.52 (s, 1H), 5.56–5.49 (m, 1H), 4.80–4.68 (m, 2H), 3.40–3.28 (m, 1H), 2.97 (dd, 1H, *J* = 6.9 and 15.6 Hz), 2.42–2.31 (m, 2H), 2.27–2.13 (m, 4H), 1.84–1.62 (m, 4H). ¹³C NMR (100 MHz, CDCl₃) δ: 180.5, 175.2, 168.9, 149.9, 134.7, 132.3, 130.6, 129.7, 127.0, 125.7, 124.4, 119.2, 114.8, 84.6, 53.2, 30.9, 29.6, 26.3, 25.3, 22.4, 22.1. ESI-MS: (*m/z*) [C₂₁H₂₁N₃O₃][−]: 362.1893. Cald. for [C₂₁H₂₁N₃O₃][−]: 362.1510.

2-((4-(1-Hydroxycyclohexyl)-1H-1,2,3-triazol-1-yl)methyl)-2,3-dihydronaphtho[1,2-*b*]furan-4,5-dione (26). Using 1-ethynylcyclohexanol (136 mg, 1.1 mmol), **26** was obtained as a red solid (345 mg, 0.91 mmol, 90% yield); mp 92–93 °C. ¹H NMR (400 MHz, CDCl₃) δ: 8.03 (d, 1H, *J* = 7.2 Hz), 7.71–7.64 (m, 2H), 7.61–7.58 (m, 2H), 5.56–5.50 (m, 1H), 4.82–4.69 (m, 2H), 3.36–3.30 (m, 1H), 2.98 (dd, 1H, *J* = 6.7 and 15.7 Hz), 2.17 (s, 1H), 2.02–1.92 (m, 2H), 1.89–1.78 (m, 2H), 1.76–1.68 (m, 2H), 1.54–1.47 (m, 2H), 1.36–1.24 (m, 2H). ¹³C NMR (100 MHz, CDCl₃) δ: 180.4, 175.1, 168.9, 134.8, 132.3, 130.5, 129.7, 126.9, 124.5, 120.9, 114.8, 84.5, 69.6, 53.4, 38.1, 38.0, 29.6, 25.3, 22.0. ESI-MS: (*m/z*) [C₂₁H₂₁N₃O₄Na]⁺: 402.1518. Cald. for [C₂₁H₂₁N₃O₄Na]⁺: 402.1424.

2-(((4-(Tetrahydro-2H-pyran-2-yl)oxy)methyl)-1H-1,2,3-triazol-1-yl)methyl)-2,3-dihydronaphtho[1,2-*b*]furan-4,5-dione (27). Using 2-(prop-2-yn-1-yloxy)tetrahydro-2H-pyran (154 mg, 1.1 mmol), **27** was obtained as a red solid (324 mg, 0.82 mmol, 82% yield); mp 86–87 °C. ¹H NMR (400 MHz, CDCl₃) δ: 8.02 (d, 1H, *J* = 7.1 Hz), 7.72 (s, 1H), 7.67–7.63 (m, 1H), 7.59–7.56 (m, 2H), 5.56–5.49 (m, 1H), 4.89–4.64 (m, 6H), 3.90–3.82 (m, 1H), 3.54–3.51 (m, 1H), 3.37–3.30 (m, 1H), 2.97–2.91 (m, 1H), 1.80–1.66 (m, 2H), 1.57–1.50 (m, 3H). ¹³C NMR (100 MHz, CDCl₃) δ: 180.4, 175.2, 134.7, 132.3, 130.6, 129.7, 126.9, 124.5, 123.6, 114.8, 98.3, 98.3, 84.5, 62.5, 60.5, 53.3, 30.9, 30.4, 29.6, 25.3, 19.4. ESI-MS: (*m/z*) [C₂₁H₂₁N₃O₅Na]⁺: 418.1016. Cald. for [C₂₁H₂₁N₃O₅Na]⁺: 418.1379.

2-((4-(Cyclohex-1-en-1-yl)-1H-1,2,3-triazol-1-yl)methyl)-2,3-dihydronaphtho[1,2-*b*]furan-4,5-dione (28). Using 1-ethynylcyclohex-1-ene (116 mg, 1.1 mmol), **28** was obtained as a red solid (336 mg, 0.93 mmol, 93% yield); mp 112–113 °C. ¹H NMR (400 MHz, CDCl₃) δ: 8.03 (d, 1H, *J* = 7.4 Hz), 7.70–7.66 (t, 1H, *J* = 16.3 Hz), 7.61–7.54 (m, 2H), 7.54 (s, 1H), 6.55–6.49 (m,

1H), 5.56–5.49 (m, 1H), 4.80–4.68 (m, 2H), 3.33 (dd, 1H, $J = 10.2$ and 15.7 Hz), 2.96 (dd, 1H, $J = 7.0$ and 15.7 Hz), 2.41–2.16 (m, 3H), 1.82–1.61 (m, 4H), 0.92–0.83 (m, 1H). ^{13}C NMR (100 MHz, CDCl_3) δ : 180.5, 175.2, 168.9, 149.9, 134.8, 132.3, 130.6, 129.7, 127.0, 125.7, 124.5, 119.2, 114.8, 84.6, 53.2, 29.6, 26.3, 25.3, 22.4, 22.1. ESI-MS: (m/z) $[\text{C}_{21}\text{H}_{20}\text{N}_3\text{O}_3]^+$: 362.1978. Calcd. for $[\text{C}_{21}\text{H}_{20}\text{N}_3\text{O}_3]^+$: 362.1499.

2-((4-Hexyl-1H-1,2,3-triazol-1-yl)methyl)-2,3-dihydronaphtho[1,2-*b*]furan-4,5-dione (29). Using oct-1-yne (121 mg, 1.1 mmol), **29** was obtained as a red solid (332 mg, 0.91 mmol, 91% yield); mp 121–123 °C. ^1H NMR (400 MHz, CDCl_3) δ : 8.04 (d, 1H, $J = 8.2$ Hz), 7.67 (t, 1H, $J = 7.4$ Hz), 7.59–7.57 (m, 2H), 7.45 (s, 1H), 5.57–5.49 (m, 1H), 4.81–4.69 (m, 2H), 3.31 (dd, 1H, $J = 10.2$ and 15.7 Hz), 2.96 (dd, 1H, $J = 6.9$ and 15.7 Hz), 2.72 (t, 2H, $J = 7.2$ Hz), 1.67–1.60 (m, 2H), 1.35–1.28 (m, 6H), 0.89–0.86 (m, 3H). ^{13}C NMR (100 MHz, CDCl_3) δ : 180.5, 175.1, 168.9, 148.9, 134.7, 132.3, 130.5, 129.7, 126.9, 124.4, 121.6, 114.8, 84.7, 53.2, 31.5, 29.6, 29.3, 28.8, 25.6, 22.5, 14.1. ESI-MS: (m/z) $[\text{C}_{21}\text{H}_{24}\text{N}_3\text{O}_3]^+$: 366.2340. Calcd. for $[\text{C}_{21}\text{H}_{24}\text{N}_3\text{O}_3]^+$: 366.1812.

General procedure for the synthesis of compounds **30**, **31** and **33–35**

To a mixture of 256 mg (1 mmol) of **24** or 269 mg (1.0 mmol) of **32** with $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ (12.48 mg, 5 mol %) and sodium ascorbate (19.81 mg, 5 mol%) in 8 mL $\text{CH}_2\text{Cl}_2/\text{H}_2\text{O}$ (1 : 1 v/v), the corresponding alkyne-carbohydrate (1.1 equivalents) was added. The mixture was stirred overnight at room temperature. The organic phase was extracted with dichloromethane, dried with NaSO_4 and concentrated under reduced pressure. The residue obtained was purified by column chromatography on silica gel using as eluent a gradient mixture of hexane/ethyl acetate with increasing polarity. The products **33–35** were obtained as diastereomers (diastereomer ratio 1 : 1).

(2S,3S,4R,5S,6S)-2-(Acetoxymethyl)-6-((1-((4,5-dioxo-2,3,4,5-tetrahydronaphtho[1,2-*b*]furan-2-yl)methyl)-1H-1,2,3-triazol-4-yl)methoxy)tetrahydro-2H-pyran-3,4,5-triyl triacetate (30). Using (2*R*,3*R*,4*S*,5*R*,6*R*)-2-(acetoxymethyl)-6-(prop-2-yn-1-yloxy)tetrahydro-2*H*-pyran-3,4,5-triyl triacetate (424 mg, 1.1 mmol), **30** was obtained as a yellow solid (544 mg, 0.8 mmol, 85% yield); mp 117–119 °C. ^1H NMR (400 MHz, CDCl_3) δ : 8.06 (bs, 1H), 7.76–7.67 (m, 2H), 7.61–7.59 (bs, 2H), 5.53–5.48 (m, 1H), 5.25–4.70 (m, 8H), 4.25–4.15 (m, 2H), 3.77 (d, 1H, $J = 4$ Hz), 3.38–3.32 (m, 1H), 2.98–2.93 (m, 1H), 2.08 (s, 3H), 2.04 (s, 3H), 2.01 (s, 3H), 1.97 (s, 3H). ^{13}C NMR (100 MHz, CDCl_3) δ : 180.4, 178.7, 175.2, 173.8, 170.6, 170.1, 169.4, 144.7, 134.9, 132.4, 130.5, 129.7, 126.8, 124.6, 123.8, 114.7, 100.0, 84.5, 72.7, 72.0, 71.2, 68.3, 63.2, 61.8, 53.4, 29.7, 20.7, 20.7, 20.6, 20.6. ESI-MS: (m/z) $[\text{C}_{30}\text{H}_{31}\text{N}_3\text{O}_{13}\text{Na}]^+$: 664.1768. Calcd. for $[\text{C}_{30}\text{H}_{31}\text{N}_3\text{O}_{13}\text{Na}]^+$: 664.1755.

(2R,3S,4S,5R,6R)-2-(Acetoxymethyl)-6-((1-((4,5-dioxo-2,3,4,5-tetrahydronaphtho[1,2-*b*]furan-2-yl)methyl)-1H-1,2,3-triazol-4-yl)methoxy)tetrahydro-2H-pyran-3,4,5-triyl triacetate (31). Using (2*R*,3*S*,4*S*,5*R*,6*R*)-2-(acetoxymethyl)-6-(prop-2-yn-1-yloxy)tetrahydro-2*H*-pyran-3,4,5-triyl triacetate, **31** was obtained as a yellow solid (512 mg, 0.8 mmol, 80% yield); mp 121–122 °C. ^1H NMR (400 MHz, CDCl_3) δ : 8.09 (bs, 1H), 7.73–7.69 (m, 2H), 7.63–7.58

(m, 2H), 5.55–5.44 (m, 1H), 5.39 (s, 1H), 5.26–5.15 (m, 1H), 5.06–4.95 (m, 2H), 4.81–4.67 (m, 4H), 4.20–4.12 (m, 2H), 3.98 (s, 1H), 3.39–3.33 (m, 1H), 2.98–2.93 (m, 1H), 2.15 (s, 3H), 2.06 (s, 3H), 1.99 (s, 6H). ^{13}C NMR (100 MHz, CDCl_3) δ : 180.4, 175.3, 171.1, 170.1, 170.0, 169.6, 169.5, 144.8, 134.9, 132.4, 130.6, 129.8, 126.9, 124.6, 123.6, 114.7, 100.0, 84.6, 70.9, 70.8, 68.8, 67.1, 63.0, 61.2, 60.4, 53.1, 29.7, 21.0, 20.6, 20.5, 19.0, 14.2. ESI-MS: (m/z) $[\text{C}_{30}\text{H}_{31}\text{N}_3\text{O}_{13}\text{Na}]^+$: 664.1762. Calcd. for $[\text{C}_{30}\text{H}_{31}\text{N}_3\text{O}_{13}\text{Na}]^+$: 664.1755.

(2R,3R,4S,5R,6R)-2-(Acetoxymethyl)-6-((1-(2,2-dimethyl-4,5-dioxo-2,3,4,5-tetrahydronaphtho[1,2-*b*]furan-3-yl)-1H-1,2,3-triazol-4-yl)methoxy)tetrahydro-2H-pyran-3,4,5-triyl triacetate (33). Using (2*R*,3*R*,4*S*,5*R*,6*R*)-2-(acetoxymethyl)-6-(prop-2-yn-1-yloxy)tetrahydro-2*H*-pyran-3,4,5-triyl triacetate (424 mg, 1.1 mmol), **33** was obtained as a yellow solid (537 mg, 0.82 mmol, 82% yield); mp 88–90 °C. ^1H NMR (400 MHz, CDCl_3) δ : 8.20–8.17 (m, 1H), 7.82–7.71 (m, 3H), 7.55/7.51 (s, 1H), 5.98/5.97 (s, 1H), 5.25–5.15 (m, 1H), 5.07 (t, 1H, $J = 10$ Hz), 5.00–4.78 (m, 3H), 4.67 (t, 1H, $J = 7.6$ Hz), 4.30–4.10 (m, 2H), 3.77–3.70 (m, 1H), 2.07, 2.01, 2.00, 1.98, 1.97, 1.77 (six singlets observed, 12H), 1.19 (s, 3H), 0.98–0.83 (m, 3H). ^{13}C NMR (100 MHz, CDCl_3) δ : 180.0, 180.0, 174.6, 174.5, 171.3, 171.2, 170.7, 170.6, 170.1, 170.1, 169.6, 169.5, 169.4, 144.4, 144.2, 134.8, 134.8, 133.4, 133.4, 131.6, 131.5, 130.0, 130.0, 126.6, 126.6, 125.6, 125.5, 122.6, 122.3, 111.1, 100.2, 99.9, 95.9, 95.8, 72.8, 72.6, 71.9, 71.3, 71.2, 68.3, 68.3, 67.0, 66.9, 63.3, 63.0, 61.8, 61.7, 31.5, 29.0, 27.7, 27.6, 22.6, 22.6, 21.1, 20.7, 20.6, 20.5, 20.5, 20.4, 14.1, 11.4. ESI-MS: (m/z) $[\text{C}_{31}\text{H}_{33}\text{N}_3\text{O}_{13}\text{Na}]^+$: 678.1925. Calcd. for $[\text{C}_{31}\text{H}_{33}\text{N}_3\text{O}_{13}\text{Na}]^+$: 678.1911.

(2R,3S,4S,5R,6R)-2-(Acetoxymethyl)-6-((1-(2,2-dimethyl-4,5-dioxo-2,3,4,5-tetrahydronaphtho[1,2-*b*]furan-3-yl)-1H-1,2,3-triazol-4-yl)methoxy)tetrahydro-2H-pyran-3,4,5-triyl triacetate (34). Using (2*R*,3*S*,4*S*,5*R*,6*R*)-2-(acetoxymethyl)-6-(prop-2-yn-1-yloxy)tetrahydro-2*H*-pyran-3,4,5-triyl triacetate (424 mg, 1.1 mmol), **34** was obtained as a yellow solid (543 mg, 0.83 mmol, 83% yield); mp 118–119 °C. ^1H NMR (400 MHz, CDCl_3) δ : 8.20/8.18 (s, 1H), 7.88–7.78 (m, 3H), 7.60 (s, 1H), 5.99/5.98 (s, 1H), 5.33–5.20 (m, 3H), 4.95–4.80 (m, 2H), 4.77–4.66 (m, 1H), 4.3 (dd, 1H, $J = 12$ and 4.8 Hz), 4.18–4.00 (m, 2H), 2.09, 1.97, 1.95, 1.78 (four singlets observed, 12H), 1.24/1.23 (s, 3H), 0.90–0.84 (m, 3H). ^{13}C NMR (100 MHz, CDCl_3) δ : 180.0, 174.6, 171.4, 171.3, 170.7, 170.2, 170.0, 170.0, 169.7, 143.6, 134.8, 134.7, 133.4, 131.6, 131.6, 130.0, 129.9, 126.7, 126.6, 125.7, 122.4, 122.3, 111.1, 111.1, 97.1, 96.8, 95.9, 95.9, 69.5, 69.5, 69.0, 68.8, 68.7, 67.0, 67.0, 66.1, 66.0, 62.4, 62.3, 61.4, 61.3, 36.1, 34.6, 31.6, 29.0, 27.7, 27.7, 22.6, 22.6, 22.1, 21.3, 20.8, 20.7, 20.6, 20.4, 18.7, 14.1, 11.4. ESI-MS: (m/z) $[\text{C}_{31}\text{H}_{33}\text{N}_3\text{O}_{13}\text{Na}]^+$: 678.1913. Calcd. for $[\text{C}_{31}\text{H}_{33}\text{N}_3\text{O}_{13}\text{Na}]^+$: 678.1911.

(2R,3R,4S,5S,6S)-2-(Acetoxymethyl)-6-((1-(2,2-dimethyl-4,5-dioxo-2,3,4,5-tetrahydronaphtho[1,2-*b*]furan-3-yl)-1H-1,2,3-triazol-4-yl)methoxy)tetrahydro-2H-pyran-3,4,5-triyl triacetate (35). Using (2*R*,3*R*,4*S*,5*S*,6*S*)-2-(acetoxymethyl)-6-(prop-2-yn-1-yloxy)tetrahydro-2*H*-pyran-3,4,5-triyl triacetate (424 mg, 1.1 mmol), **35** was obtained as a yellow solid (556 mg, 0.85 mmol, 85% yield); mp 104–107 °C. ^1H NMR (400 MHz, CDCl_3) δ : 8.20/8.18 (m, 1H), 7.84–7.72 (m, 3H), 7.52/7.49 (s, 1H), 5.98/5.95 (s, 1H), 5.39/5.38 (s, 1H), 5.22–5.12 (m, 1H), 5.03–4.92 (m, 2H),

4.84–4.77 (m, 1H), 4.65–4.62 (m, 1H), 4.15–4.12 (m, 2H), 3.98–3.93 (m, 1H), 2.13, 2.11, 2.04, 2.03, 1.97, 1.96, 1.77 (seven singlets observed, 12H), 1.20/1.19 (s, 3H), 0.90–0.83 (m, 3H). ^{13}C NMR (100 MHz, CDCl_3) δ : 180.0, 180.0, 174.5, 174.5, 171.2, 170.4, 170.2, 170.0, 170.0, 169.8, 169.8, 169.6, 169.6, 144.5, 144.2, 134.8, 133.4, 131.5, 130.0, 126.6, 125.5, 125.5, 122.5, 122.1, 111.3, 111.1, 100.7, 100.3, 96.0, 95.9, 70.9, 70.8, 70.7, 68.8, 68.7, 67.1, 67.0, 66.9, 66.8, 63.2, 62.7, 61.3, 61.3, 31.6, 29.0, 27.7, 22.6, 22.6, 21.2, 21.1, 20.7, 20.6, 20.5, 14.1, 11.4. ESI-MS: (m/z) [$\text{C}_{31}\text{H}_{33}\text{N}_3\text{O}_{13}\text{Na}$] $^+$: 678.1938. Cald. for [$\text{C}_{31}\text{H}_{33}\text{N}_3\text{O}_{13}\text{Na}$] $^+$: 678.1911.

Acknowledgements

This research was funded by grants from the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq 480719/2012-8, PVE 401193/2014-4, PQ-SR CNPq 302613/2013-7), FAPEMIG (Edital 01/2014), CAPES, FAPERJ, FAPEAL, CAPES/PROCAD and PAPES/Fiocruz.

Notes and references

- (a) WHO, Chagas disease in Latin America: an epidemiological update based on 2010 estimates, *Wkly. Epidemiol. Rec.*, 2015, **90**, 33; (b) A. Rassi Júnior, A. Rassi and J. M. de Rezende, *Infect. Dis. Clin.*, 2012, **26**, 275.
- M. Steindel, L. K. Pacheco, D. Scholl, M. Soares, M. H. de Moraes, I. Eger, C. Kosmann, T. C. M. Sincero, P. H. Stoco, S. M. F. Murta, C. J. Carvalho-Pinto and E. C. Grisard, *Diagn. Microbiol. Infect. Dis.*, 2008, **60**, 25.
- J. C. P. Dias, V. A. Neto and E. J. A. Luna, *Rev. Soc. Bras. Med. Trop.*, 2011, **44**, 375.
- J. R. Coura and J. C. P. Dias, *Mem. Inst. Oswaldo Cruz*, 2009, **104**, 31.
- J. Gascon, C. Bern and M. J. Pinazo, *Acta Trop.*, 2010, **115**, 22.
- G. A. Schmunis and Z. E. Yadon, *Acta Trop.*, 2010, **115**, 14.
- A. Rassi Júnior, A. Rassi and J. A. Marin-Neto, *Lancet*, 2010, **375**, 1388.
- W. de Souza, *Curr. Pharm. Des.*, 2002, **8**, 269.
- M. N. Soeiro and S. L. de Castro, *Open Med. Chem. J.*, 2011, **5**, 21.
- (a) K. W. Wellington, *RSC Adv.*, 2015, **5**, 20309; (b) J. A. Urbina, *J. Eukaryotic Microbiol.*, 2015, **62**, 149.
- G. Powis, *Pharmacol. Ther.*, 1987, **35**, 57.
- E. A. Hillard, F. C. Abreu, D. C. M. Ferreira, G. Jaouen, M. O. F. Goulart and C. Amatore, *Chem. Commun.*, 2008, 2612.
- B. Hazra, M. Das Sarma and U. Sanyal, *J. Chromatogr. B: Anal. Technol. Biomed. Life Sci.*, 2004, **812**, 259.
- C. C. Nawrat and C. J. Moody, *Angew. Chem., Int. Ed.*, 2014, **53**, 2056.
- (a) E. N. da Silva Júnior, G. A. M. Jardim, R. F. S. Menna-Barreto and S. L. de Castro, *J. Braz. Chem. Soc.*, 2014, **25**, 1780; (b) A. Cavalli and M. L. Bolognesi, *J. Med. Chem.*, 2009, **52**, 7339; (c) K. Vázquez, C. Espinosa-Bustos, J. Soto-Delgado, R. A. Tapia, J. Varela, E. Birriel, R. Segura, J. Pizarro, H. Cerecetto, M. González, M. Paulino and C. O. Salas, *RSC Adv.*, 2015, **5**, 65153.
- R. Kharb, P. C. Sharma and M. S. Yar, *J. Enzyme Inhib. Med. Chem.*, 2011, **26**, 1.
- S. Scodavolpe, S. Quaranta, B. Lacarelle and C. Solas, *Ann. Biol. Clin.*, 2014, **72**, 391.
- R. Huisgen, G. Szeimies and L. Möbius, *Chem. Ber.*, 1967, **100**, 2494.
- (a) H. C. Kolb and K. B. Sharpless, *Drug Discovery Today*, 2003, **8**, 1128; (b) H. C. Kolb, M. G. Finn and K. B. Sharpless, *Angew. Chem., Int. Ed.*, 2001, **40**, 2004; (c) C. W. Tornøe, C. Christensen and M. Meldal, *J. Org. Chem.*, 2002, **67**, 3057; (d) V. V. Rostovtsev, L. G. Green, V. V. Fokin and K. B. Sharpless, *Angew. Chem., Int. Ed.*, 2002, **41**, 2596; (e) M. Meldal and C. W. Tornøe, *Chem. Rev.*, 2008, **108**, 2952.
- (a) S. G. Agalave, S. R. Maujan and V. S. Pore, *Chem.-Asian J.*, 2011, **6**, 2696; (b) J. Hou, X. Liu, J. Shen, G. Zhao and P. G. Wang, *Expert Opin. Drug Discovery*, 2012, **7**, 489; (c) D. Pasini, *Molecules*, 2013, **18**, 9512.
- S. L. de Castro, F. S. Emery and E. N. da Silva Júnior, *Eur. J. Med. Chem.*, 2013, **69**, 678.
- A. V. Pinto and S. L. de Castro, *Molecules*, 2009, **14**, 4570.
- (a) K. C. G. Moura, K. Salomão, R. F. S. Menna-Barreto, F. S. Emery, M. C. F. R. Pinto, A. V. Pinto and S. L. de Castro, *Eur. J. Med. Chem.*, 2004, **39**, 639; (b) K. C. G. Moura, F. S. Emery, C. N. Pinto, M. C. F. R. Pinto, A. P. Dantas, K. Salomão, S. L. de Castro and A. V. Pinto, *J. Braz. Chem. Soc.*, 2001, **12**, 325; (c) A. V. Pinto, C. Neves-Pinto, M. C. F. R. Pinto, R. M. Santa Rita, C. Pezzella and S. L. de Castro, *Arzneim.-Forsch.*, 1997, **47**, 74; (d) C. Neves-Pinto, A. P. Dantas, K. C. G. Moura, F. S. Emery, P. F. Polequevitch, M. C. F. R. Pinto, S. L. de Castro and A. V. Pinto, *Arzneim.-Forsch.*, 2000, **50**, 1120.
- (a) R. F. S. Menna-Barreto, J. R. Corrêa, C. M. Cascabulho, M. C. Fernandes, A. V. Pinto, M. J. Soares and S. L. de Castro, *Parasitology*, 2009, **136**, 499; (b) R. F. S. Menna-Barreto, J. R. Corrêa, A. V. Pinto, M. J. Soares and S. L. de Castro, *Parasitol. Res.*, 2007, **101**, 895; (c) R. F. S. Menna-Barreto, A. Henriques-Pons, A. V. Pinto, J. A. Morgado-Diaz, M. J. Soares and S. L. de Castro, *J. Antimicrob. Chemother.*, 2005, **56**, 1034.
- E. N. da Silva Júnior, M. C. B. V. de Souza, M. C. Fernandes, R. F. S. Menna-Barreto, M. C. F. R. Pinto, F. A. Lopes, C. A. de Simone, C. K. Z. Andrade, A. V. Pinto, V. F. Ferreira and S. L. de Castro, *Bioorg. Med. Chem.*, 2008, **16**, 5030.
- E. N. da Silva Júnior, R. F. S. Menna-Barreto, M. C. F. R. Pinto, R. S. F. Silva, D. V. Teixeira, M. C. B. V. de Souza, C. A. de Simone, S. L. de Castro, V. F. Ferreira and A. V. Pinto, *Eur. J. Med. Chem.*, 2008, **42**, 1774.
- E. N. da Silva Júnior, I. M. M. de Melo, E. B. T. Diogo, V. A. Costa, J. D. de Souza Filho, W. O. Valença, C. A. Camara, R. N. de Oliveira, A. S. de Araújo, F. S. Emery, M. R. dos Santos, C. A. de Simone, R. F. S. Menna-Barreto and S. L. de Castro, *Eur. J. Med. Chem.*, 2012, **52**, 304.

- 28 R. S. F. Silva, E. M. Costa, U. L. T. Trindade, D. V. Teixeira, M. C. F. R. Pinto, G. L. Santos, V. R. S. Malta, C. A. de Simone, A. V. Pinto and S. L. de Castro, *Eur. J. Med. Chem.*, 2006, **41**, 526.
- 29 R. F. S. Menna-Barreto, R. L. S. Gonçalves, E. M. Costa, R. S. F. Silva, A. V. Pinto, M. F. Oliveira and S. L. de Castro, *Free Radical Biol. Med.*, 2009, **47**, 644.
- 30 A. Dondoni, *Chem.-Asian J.*, 2007, **2**, 700.
- 31 I. Carvalho, P. Andrade, V. L. Campo, P. M. Guedes, R. Sesti-Costa, J. S. Silva, S. Schenkman, S. Dedola, L. Hill, M. Rejzek, S. A. Nepogodiev and R. A. Field, *Bioorg. Med. Chem.*, 2010, **18**, 2412.
- 32 V. L. Campo, R. Sesti-Costa, Z. A. Carneiro, J. S. Silva, S. Schenkman and I. Carvalho, *Bioorg. Med. Chem.*, 2012, **20**, 145.
- 33 P. Singh, A. Anand and V. Kumar, *Eur. J. Med. Chem.*, 2014, **85**, 758.
- 34 (a) Z. Nowakowska, *Eur. J. Med. Chem.*, 2007, **42**, 125; (b) D. I. Batovska and I. T. Todorova, *Curr. Clin. Pharmacol.*, 2010, **5**, 1; (c) E. H. Zhang, R. F. Wang, S. Z. Guo and B. Liu, *J. Evidence-Based Complementary Altern. Med.*, 2013, **2013**, 815621; (d) M. J. Matos, S. Vazquez-Rodriguez, E. Uriarte and L. Santana, *Expert Opin. Ther. Pat.*, 2015, **25**, 351.
- 35 M. Roussaki, B. Hall, S. C. Lima, A. C. da Silva, S. Wilkinson and A. Detsi, *Bioorg. Med. Chem. Lett.*, 2013, **23**, 6436.
- 36 J. Ryczak, M. Papini, A. Lader, A. Nasereddin, D. Kopelyanskiy, L. Preu, C. L. Jaffe and C. Kunick, *Eur. J. Med. Chem.*, 2013, **64**, 396.
- 37 Z. Qiao, Q. Wang, F. Zhang, Z. Wang, T. Bowling, B. Nare, R. T. Jacobs, J. Zhang, D. Ding, Y. Liu and H. Zhou, *J. Med. Chem.*, 2012, **55**, 3553.
- 38 C. Ibis, A. F. Tuyun, H. Bahar, S. S. Ayla, M. V. Stasevych, R. Y. Musyanovych, O. Komarovska-Porokhnyavets and V. Novikov, *Med. Chem. Res.*, 2013, **22**, 2879.
- 39 V. Jamier, L. A. Ba and C. Jacob, *Chem.-Eur. J.*, 2010, **16**, 10920.
- 40 (a) S. Shaaban, R. Diestel, B. Hinkelmann, Y. Muthukumar, R. P. Verma, F. Sasse and C. Jacob, *Eur. J. Med. Chem.*, 2012, **58**, 192; (b) F. H. Fry, A. L. Holme, N. M. Giles, G. I. Giles, C. Collins, K. Holt, S. Pariagh, T. Gelbrich, M. B. Hursthouse, N. J. Gutowski and C. Jacob, *Org. Biomol. Chem.*, 2005, **3**, 2579; (c) M. Doering, B. Diesel, M. C. H. Gruhlke, U. M. Viswanathan, D. Manikov, M. Chovanec, T. Burkholz, A. J. Slusarenko, A. K. Kiemer and C. Jacob, *Tetrahedron*, 2012, **68**, 10577.
- 41 A. V. Pinto, M. C. F. R. Pinto and C. G. T. de Oliveira, *An. Acad. Bras. Cienc.*, 1982, **54**, 108.
- 42 E. H. G. Cruz, C. M. B. Hussene, G. G. Dias, E. B. T. Diogo, I. M. M. de Melo, B. L. Rodrigues, M. G. da Silva, W. O. Valença, C. A. Camara, R. N. de Oliveira, Y. G. de Paiva, M. O. F. Goulart, B. C. Cavalcanti, C. Pessoa and E. N. da Silva Júnior, *Bioorg. Med. Chem.*, 2014, **22**, 1608.
- 43 (a) R. T. Blickenstaff, W. R. Hanson, S. Reddy and R. Witt, *Bioorg. Med. Chem.*, 1995, **3**, 917; (b) V. Tomar, G. Bhattacharjee, Kamaluddin, S. Rajakumar, K. Srivastava and S. K. Puri, *Eur. J. Med. Chem.*, 2010, **45**, 745; (c) F. L. Ansari, S. Umbreen, L. Hussain, T. Makhmoor, S. A. Nawaz, M. A. Lodhi, S. N. Khan, F. Shaheen, M. I. Choudhary and Atta-ur-Rahman, *Chem. Biodiversity*, 2005, **2**, 487.
- 44 L. F. Fieser and M. Fieser, *J. Am. Chem. Soc.*, 1948, **70**, 3215.
- 45 G. A. M. Jardim, T. T. Guimarães, M. C. F. R. Pinto, B. C. Cavalcanti, K. M. de Farias, C. Pessoa, C. C. Gatto, D. K. Nair, I. N. N. Namboothiri and E. N. da Silva Júnior, *Med. Chem. Commun.*, 2015, **6**, 120.
- 46 A. J. M. da Silva, C. D. Buarque, F. V. Brito, L. Aurelian, L. F. Macedo, L. H. Malkas, R. J. Hickey, D. V. S. Lopes, F. Noël, Y. L. B. Murakami, N. M. V. Silva, P. A. Melo, R. R. B. Caruso, N. G. Castro and P. R. R. Costa, *Bioorg. Med. Chem.*, 2002, **10**, 2731.
- 47 (a) G. A. M. Jardim, E. H. G. da Cruz, W. O. Valença, J. M. Resende, B. L. Rodrigues, D. F. Ramos, R. N. de Oliveira, P. E. A. Silva and E. N. da Silva Júnior, *J. Braz. Chem. Soc.*, 2015, **26**, 1013; (b) I. C. Chipoline, V. F. Ferreira and F. C. da Silva, *Curr. Org. Synth.*, 2015, **12**, 565.
- 48 Z. Wang, M. Matin and S. Sheikh, *Molecules*, 2005, **10**, 1325.
- 49 K. P. R. Kartha and R. A. Field, *Tetrahedron*, 1997, **53**, 11753.
- 50 R. Yang, H. Ding, Y. Song, W. Xiao, Q. Xiao and J. Wu, *Lett. Org. Chem.*, 2008, **5**, 518.
- 51 A. A. Vieira, I. R. Brandão, W. O. Valença, C. A. de Simone, B. C. Cavalcanti, C. Pessoa, T. R. Carneiro, A. L. Braga and E. N. da Silva Júnior, *Eur. J. Med. Chem.*, 2015, **101**, 254.
- 52 E. N. da Silva Júnior, T. T. Guimarães, R. F. S. Menna-Barreto, M. C. F. R. Pinto, C. A. de Simone, C. Pessoa, B. C. Cavalcanti, J. R. Sabino, C. K. Z. Andrade, M. O. F. Goulart, S. L. de Castro and A. V. Pinto, *Bioorg. Med. Chem.*, 2010, **18**, 3224.
- 53 M. C. Fernandes, E. N. da Silva Júnior, A. V. Pinto, S. L. de Castro and R. F. S. Menna-Barreto, *Parasitology*, 2012, **139**, 26.
- 54 Y. G. de Paiva, F. R. Ferreira, T. L. Silva, E. Labbé, O. Buriez, C. Amatore and M. O. F. Goulart, *Curr. Top. Med. Chem.*, 2015, **15**, 136.
- 55 (a) M. O. F. Goulart, C. L. Zani, J. Tonholo, L. R. Freitas, F. C. de Abreu, A. B. Oliveira, D. S. Raslan, S. Starling and E. Chiari, *Bioorg. Med. Chem. Lett.*, 1997, **7**, 2043; (b) E. N. da Silva Júnior, M. A. B. F. de Moura, A. V. Pinto, M. C. F. R. Pinto, M. C. B. V. Souza, A. J. Araújo, C. Pessoa, L. V. Costa-Lotufó, R. Montenegro, M. O. de Moraes, M. O. F. Goulart and V. F. Ferreira, *J. Braz. Chem. Soc.*, 2009, **20**, 635.
- 56 E. B. T. Diogo, G. G. Dias, B. L. Rodrigues, T. T. Guimarães, W. O. Valença, C. A. Camara, R. N. de Oliveira, M. G. da Silva, V. F. Ferreira, Y. G. de Paiva, M. O. F. Goulart, R. F. S. Menna-Barreto, S. L. de Castro and E. N. da Silva Júnior, *Bioorg. Med. Chem.*, 2013, **21**, 6337.
- 57 G. I. Giles, K. M. Tasker, R. J. K. Johnson, C. Jacob, C. Peers and K. N. Green, *Chem. Commun.*, 2001, 2490.
- 58 H. Viertler, J. Gruber and V. L. Pardini, Anodic oxidation of sulfur- and selenium-containing compounds, in *Organic Electrochemistry*, ed. H. Lund and O. Hammerich, Marcel

- Dekker, Inc., New York, 4th edn, 2001, ch. 17, H. Lund and M. M. Baizer, pp. 621–668 and included references.
- 59 R. R. France, N. V. Rees, J. D. Wadhawan, A. J. Fairbanks and R. G. Compton, *Org. Biomol. Chem.*, 2004, **2**, 2188.
- 60 F. R. Ferreira, S. B. Ferreira, A. J. Araújo, J. D. B. M. Filho, C. Pessoa, M. O. Moraes, L. V. Costa-Lotufo, R. C. Montenegro, F. C. da Silva, V. F. Ferreira, J. G. da Costa, F. C. de Abreu and M. O. F. Goulart, *Electrochim. Acta*, 2013, **110**, 634.
- 61 B. Dakova, L. Lambews and M. Evers, *Electrochim. Acta*, 1994, **39**, 2363.
- 62 M. O. F. Goulart, F. C. Abreu and A. M. O. Brett, *Electroanalysis*, 2002, **14**, 29.
- 63 S. L. de Castro and M. N. L. Meirelles, *Mem. Inst. Oswaldo Cruz*, 1990, **85**, 95.