

The Essential Role of Cholesterol Metabolism in the Intracellular Survival of *Mycobacterium leprae* Is Not Coupled to Central Carbon Metabolism and Energy Production

Maria Angela M. Marques,^{a,b} Marcia Berrêdo-Pinho,^a Thabatta L. S. A. Rosa,^a Venugopal Pujari,^b Robertha M. R. Lemes,^a Leticia M. S. Lery,^a Carlos Adriano M. Silva,^b Ana Carolina R. Guimarães,^c Georgia C. Atella,^d William H. Wheat,^b Patrick J. Brennan,^b Dean C. Crick,^b John T. Belisle,^b Maria Cristina V. Pessolani^a

Laboratório de Microbiologia Celular, Instituto Oswaldo Cruz, Fundação Oswaldo Cruz, Rio de Janeiro, RJ, Brazil^a; Department of Microbiology, Immunology and Pathology, Colorado State University, Fort Collins, Colorado, USA^b; Laboratório de Genômica Funcional e Bioinformática, Instituto Oswaldo Cruz, Fundação Oswaldo Cruz, Rio de Janeiro, RJ, Brazil^c; Laboratório de Bioquímica de Lipídeos e Lipoproteínas, Instituto de Bioquímica Médica, Universidade Federal do Rio de Janeiro, Rio de Janeiro, RJ, Brazil^d

ABSTRACT

Mycobacterium leprae induces the formation of lipid droplets, which are recruited to pathogen-containing phagosomes in infected macrophages and Schwann cells. Cholesterol is among the lipids with increased abundance in *M. leprae*-infected cells, and intracellular survival relies on cholesterol accumulation. The present study investigated the capacity of *M. leprae* to acquire and metabolize cholesterol. *In silico* analyses showed that oxidation of cholesterol to cholest-4-en-3-one (cholestenone), the first step of cholesterol degradation catalyzed by the enzyme 3 β -hydroxysteroid dehydrogenase (3 β -HSD), is apparently the only portion of the cholesterol catabolic pathway seen in *Mycobacterium tuberculosis* preserved by *M. leprae*. Incubation of bacteria with radiolabeled cholesterol confirmed the *in silico* predictions. Radiorespirometry and lipid analyses performed after incubating *M. leprae* with [4-¹⁴C]cholesterol or [26-¹⁴C]cholesterol showed the inability of this pathogen to metabolize the sterol rings or the side chain of cholesterol as a source of energy and carbon. However, the bacteria avidly incorporated cholesterol and, as expected, converted it to cholestenone both *in vitro* and *in vivo*. Our data indicate that *M. leprae* has lost the capacity to degrade and utilize cholesterol as a nutritional source but retains the enzyme responsible for its oxidation to cholestenone. Thus, the essential role of cholesterol metabolism in the intracellular survival of *M. leprae* is uncoupled from central carbon metabolism and energy production. Further elucidation of cholesterol metabolism in the host cell during *M. leprae* infection will establish the mechanism by which this lipid supports *M. leprae* intracellular survival and will open new avenues for novel leprosy therapies.

IMPORTANCE

Our study focused on the obligate intracellular pathogen *Mycobacterium leprae* and its capacity to metabolize cholesterol. The data make an important contribution for those interested in understanding the mechanisms of mycobacterial pathogenesis, since they indicate that the essential role of cholesterol for *M. leprae* intracellular survival does not rely on its utilization as a nutritional source. Our findings reinforce the complexity of cholesterol's role in sustaining *M. leprae* infection. Further elucidation of cholesterol metabolism in the host cell during *M. leprae* infection will establish the mechanism by which this lipid supports *M. leprae* intracellular survival and will open new avenues for novel leprosy therapies.

Leprosy is a granulomatous disease caused by *Mycobacterium leprae*, an obligate intracellular organism that affects mainly the skin and peripheral nerves. This ancient disease, which can be considered one of the oldest human infections (1), is still a significant public health problem in developing countries, including India, Brazil, and Indonesia, which collectively account for 83% of all new cases worldwide (2). Although the global prevalence of leprosy has declined from 5.2 million in the 1980s to approximately 200,000 today following the implementation of multidrug therapy (MDT) (2), the number of new cases of leprosy in regions where the disease is endemic is an indication that the transmission of *M. leprae* is still an issue. Moreover, at least in Brazil, the prevalence of undiagnosed cases in areas with a high burden of disease is reported to be much higher than the registered prevalence (3). MDT is effective in bacterial killing but does not prevent reactional episodes, the major cause of nerve injury and physical disabilities seen in individuals affected by the disease (4). Thus, novel drug combinations are needed for a better management of leprosy patients and prevention of physical sequelae.

Received 30 July 2015 Accepted 5 September 2015

Accepted manuscript posted online 21 September 2015

Citation Marques MAM, Berrêdo-Pinho M, Rosa TLA, Pujari V, Lemes RMR, Lery LMS, Silva CAM, Guimarães ACR, Atella GC, Wheat WH, Brennan PJ, Crick DC, Belisle JT, Pessolani MCV. 2015. The essential role of cholesterol metabolism in the intracellular survival of *Mycobacterium leprae* is not coupled to central carbon metabolism and energy production. *J Bacteriol* 197:3698–3707. doi:10.1128/JB.00625-15.

Editor: P. de Boer

Address correspondence to Maria Cristina V. Pessolani, cpessola@gmail.com. D.C.C., J.T.B., and M.C.V.P. are co-senior authors and contributed equally to the study.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/JB.00625-15>.

Copyright © 2015 Marques et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](http://creativecommons.org/licenses/by/4.0/).

The leprosy bacillus is an obligate intracellular pathogen with preferential tropism for macrophages of the skin and Schwann cells of the peripheral nerves. The inability of *M. leprae* to grow extracellularly or in axenic medium can be explained by the reduced coding capacity of its degenerate genome (1,605 functional genes and 1,115 pseudogenes) (5). These characteristics and a lack of experimental models of disease have hampered the study of host-pathogen interactions in leprosy. Nevertheless, insight into the pathogenesis of leprosy can be gained through the use of *in vitro* models of infection that allow dissection of host cell-pathogen interactions and complementation with *ex vivo* data based on analyses of clinical samples from leprosy patients.

As early as 1863, Virchow observed the leprosy bacillus residing inside foamy macrophages (called Virchow's cells) (6). With the discovery and characterization of *M. leprae*-derived lipids, such as phthiocerol dimycocerosate (PDIM) and phenolic glycolipid I (PGL-I), it was initially believed the lipid-heavy Virchow's cells resulted from these bacterial products (7, 8). However, recent reports demonstrate that Virchow's cells accumulate host-derived lipids (9, 10). Virchow's cells found in dermal lesions are also highly positive for adipose differentiation-related protein (ADRP), a classical cholesterol-containing lipid droplet (LD) marker. This suggests that the foamy phenotype is in part derived from the accumulation of LDs (9, 11). The ADRP marker is induced by *M. leprae* infection and localizes to *M. leprae*-containing phagosomes, indicating a close association between LDs and the pathogen-containing vacuole (11, 12). More recently, we analyzed nerve biopsy specimens and used *in vitro* cultures of Schwann cells to show that the foamy phenotype of the lepromatous leprosy (LL) nerves is also related to the capacity of *M. leprae* to induce LD biogenesis in this cell type (13, 14). Cholesterol was confirmed as a host lipid that accumulates in *M. leprae*-infected macrophages. Further, *M. leprae* increases *de novo* synthesis of cholesterol as well as exogenous uptake of low-density-lipoprotein (LDL) cholesterol by upregulating the expression of genes involved in these pathways (12). Of importance, cholesterol has been found to colocalize to *M. leprae*-containing phagosomes, and impairment of cholesterol metabolism significantly decreased intracellular bacterial survival both *in vitro* and *in vivo* (12, 15).

The dependence of mycobacteria on host lipid molecules for successful infection and persistence has been extensively analyzed in the context of *Mycobacterium tuberculosis*. *M. tuberculosis* also mediates formation of foamy cells, a process apparently critical for bacterial persistence in the host (16, 17). Additionally, a large body of literature describing the importance of cholesterol for persistence of *M. tuberculosis* in the host has accumulated (18, 19). *M. tuberculosis* possesses the ability to degrade and use cholesterol as an energy source and for the biosynthesis of mycobacterial lipids (18). This metabolic capacity for cholesterol utilization appears to be particularly important during the latent phase of *M. tuberculosis* infection, when other carbon sources become limited (18, 20). The contribution of cholesterol to the *in vivo* growth of *M. tuberculosis* and tuberculosis pathogenesis has resulted in the elucidation of *M. tuberculosis* genes directly or indirectly involved in cholesterol metabolism (for a review, see reference 21).

In light of the current understanding of cholesterol catabolism in *M. tuberculosis*, and in the face of the essential role of cholesterol for intracellular *M. leprae* survival, the present study revisited the *M. leprae* genome annotated in 2001 (5), looking for orthologous enzymes/proteins of *M. tuberculosis* genes associated with choles-

terol metabolism (21). This *in silico* analysis was complemented with investigation of the capacity of *M. leprae* to metabolize cholesterol. The data showed that *M. leprae* lost essentially all the genes associated with cholesterol catabolism in *M. tuberculosis* and consequently the capacity to degrade both the sterol rings and the lateral chain but retained the ability to oxidize cholesterol to cholest-4-en-3-one (cholestenone). These findings reinforce the complexity of cholesterol's role in sustaining *M. leprae* infection.

MATERIALS AND METHODS

Mycobacterial culture and growth medium. *M. leprae* purified from athymic *nu/nu* mouse footpads was donated by the National Hansen's Disease Program, Laboratory Research Branch, Louisiana State University, Baton Rouge, LA, USA, and by Lauro de Sousa Lima Institute, Bauru, São Paulo, Brazil. *M. tuberculosis* mc²6230 was kindly donated by William R. Jacobs, Albert Einstein College of Medicine, Bronx, NY (22). *M. tuberculosis* was grown on Middlebrook 7H9 broth supplemented with 10% oleic acid-albumin-dextrose-catalase (OADC), 0.5% glycerol, 0.05% tyloxapol, 0.2% Casamino Acids, 50 µg/ml ampicillin, and 24 µg/ml pantothenate until cell growth reached log phase. Inactivation of mycobacterial cells was achieved by heating at 100°C for 15 min.

Radiospirometry assay. *M. leprae* and *M. tuberculosis*, freshly purified from athymic *nu/nu* mouse footpads and harvested from exponential cultures, respectively, were washed three times with phosphate-buffered saline (PBS; pH 7.4) containing 0.05% tyloxapol (PBS-Ty). *M. leprae* was suspended in Middlebrook 7H9 broth supplemented with 0.1% casein hydrolysate, 0.5% BSA, 0.05% tyloxapol, 50 µg/ml ampicillin and 48 U/ml catalase. *M. tuberculosis* was suspended in Middlebrook 7H9 broth supplemented with 0.05% tyloxapol, 0.2% Casamino Acids, 50 µg/ml ampicillin, and 24 µg/ml pantothenate. The respirometry assay is based on the method of Buddemeyer (23), with minor modifications. Suspensions of mycobacterial cells (6×10^7 bacteria in 200 µl) containing 1 µCi/ml of [1-¹⁴C]palmitic acid (specific activity, 55 mCi/mmol) (American Radiolabeled Chemicals, Inc., Saint Louis, MO), [4-¹⁴C]cholesterol (specific activity, 50 mCi/mmol), or [26-¹⁴C]cholesterol (specific activity, 52 mCi/mmol) (Quotient Bioresearch Ltd., Cardiff, United Kingdom) (see cholesterol structure in Fig. 1A) were transferred to 500-µl Eppendorf tubes without caps and placed in 6-ml scintillation vials containing two filter paper strips (40 by 8 mm) impregnated with 100 µl of 2 N NaOH. After incubation for 24 and 72 h for *M. tuberculosis* and 4 days for *M. leprae*, radiolabeled CO₂ captured on the filter paper strips was measured by transferring them to a new scintillation vial containing 100 µl of concentrate acetic to neutralize the strips, adding 5 ml of Optima Gold (PerkinElmer, Waltham, MA) scintillation cocktail, and counting on a Beckman LS 600 liquid scintillation counter (Beckman Coulter, Pasadena, CA). Captured ¹⁴CO₂ was a measure of cumulative β oxidation of palmitic acid or cholesterol.

Cholesterol incorporation by *M. leprae*. Live or heat-inactivated *M. leprae* cells were incubated with [1-¹⁴C]palmitic acid, [4-¹⁴C]cholesterol, or [26-¹⁴C]cholesterol for 4 days as described in "Radiospirometry assay" above. *M. leprae* cells were collected by centrifugation at 12,000 × g for 8 min, and pellets were washed three times with PBS-Ty. The final bacterial cell pellet was suspended with 100 µl PBS, and an aliquot was transferred to a scintillation vial containing Optima Gold Plus scintillation cocktail and counted on a Beckman LS 600 liquid scintillation counter.

Lipid analysis. Mycobacterial cells labeled with [1-¹⁴C]palmitic acid, [4-¹⁴C]cholesterol, or [26-¹⁴C]cholesterol for various times were pelleted by centrifugation. The conditioned medium was also collected for extraction of lipids. The bacterial cells were washed three times with PBS-Ty. Bacterial cells and conditioned medium were extracted with chloroform-methanol (2:1) (24), and the extracts were dried under N₂ after two washes (25). Radiolabeled compounds were resolved by thin-layer chromatography (TLC) using silica gel G60 TLC plates (Millipore, Temecula, CA) developed three times in chloroform-ethyl acetate (97:3) or once in chloroform-methanol (95:5) and detected using a PhosphorImager

(Typhoon 9400 scanner; GE Amersham, Sunnyvale, CA). Lipid extracts were also spiked with 2 μg of authentic cholesterol or cholestenone (both from Sigma-Aldrich) and loaded on TLC plates, which were developed with chloroform-methanol-acetic acid (95:4:1) or chloroform-methanol-ammonium hydroxide (95:4:1). After the TLC plates had been to the phosphorimaging screen overnight, they were sprayed with a charring solution consisting of 5% phosphomolybdic acid in 100% ethanol or anisaldehyde solution (26) and heated to visualize the lipids.

Sample preparation for gas chromatography-mass spectrometry (GC-MS) and GC-MS conditions. Lipids were extracted from *M. leprae* and noninfected mouse footpad tissue as described by Bligh and Dyer (24), followed by a second extraction with chloroform-methanol (2:1) (25). The organic extracts were dried under N_2 and saponified. Briefly, 3 ml of 25% alcoholic potassium hydroxide solution (25 g of potassium hydroxide and 35 ml of sterile distilled water, brought to 100 ml with 100% ethanol) was added to the lipid pellets, mixed for 1 min, and incubated at 85°C for 1 h. The saponified material was cooled to room temperature, and the sterols were extracted by the addition of 1 ml of sterile distilled water and 3 ml of *n*-heptane, followed by vigorous mixing for 3 min. The heptane layer was collected and dried under N_2 . The isolated sterols were silylated by the addition of 50 μl bis(trimethylsilyl)trifluoroacetamide (BSTFA)-trimethylchlorosilane (TMCS) (99:1) (Sigma-Aldrich) and 50 μl pyridine and incubation for 1 h at 65°C.

GC-MS for *M. leprae* extract was performed on a Shimadzu GCMS-QP2010 Plus system (Shimadzu, Barra Funda, SP, Brazil), using an Rtx-5MS column (5% phenyl-95% dimethylpolysiloxane; 30 m by 0.25 mm by 0.25 μm ; Restex, Bellefonte, PA). The injector temperature was set at 250°C. Analytes were eluted from the column with a biphasic linear thermal gradient of 120 to 250°C at a rate of 20°C per min, followed by 250 to 280°C at a rate of 5°C per min holding at 280°C for 10 min, 280 to 330°C at a rate of 10°C per min, and holding at 330°C for 6 min. Helium was used as the carrier gas, with a linear velocity of 37.5 cm/s. The derivatized lipid extracts (5 μl) were applied to GC-MS via splitless mode injection. Electroionization was performed at -70 eV, and the quadrupole mass analyzer collected data from 30 to 500 atomic mass units (amu).

GC-MS analysis of mouse tissue extract was carried out on a Shimadzu GCMS-QP2010 Plus system, using an Agilent HP Ultra 2 column (5% phenyl-95% methylpolysiloxane; 25 m by 0.20 mm by 0.33 μm ; Agilent, Barueri, SP, Brazil). The injector temperature was set at 250°C. Analytes were eluted from the column using a thermal gradient of 50 to 270°C at a rate of 18°C per min and 270 to 300°C at a rate of 1°C per min. Helium was used as the carrier gas, with a linear velocity of 33.0 cm/s. A volume of 1 μl of sample was applied to the GC-MS. Electroionization was performed at -70 eV, and the quadrupole mass analyzer collected data from 40 to 600 amu.

To confirm chromatogram peak identities, retention indices were used. A steroid standards mix (0.5 $\mu\text{g}/\mu\text{l}$), composed of cholesterol, 7-dehydrocholesterol, ergosterol, cholestenone, α -ergosterol, stigmaterol, γ -ergosterol, lanosterol (Sigma), zymosterol, and β -sitosterol (Avanti Polar Lipids, Alabaster, AL), was processed in parallel with the sample. Additionally, the products detected by GC-MS were identified by comparing their mass spectra with those of the NIST library (NIST05).

In silico analysis. The *M. leprae* TN genome (accession number AL450380, updated 6 February 2015) and *M. tuberculosis* H37Rv genome (accession number AL123456, updated 13 June 2013) were represented using Artemis software release 16.0.0 (Sanger Institute). *M. tuberculosis* cholesterol-related proteins (reviewed in reference 21) were searched against the *M. leprae* genome using the online NCBI BLASTP tool. Proteins identified by BLAST as having more than 70% similarity and 70% coverage were considered potential matches. Those bidirectional best-hit matches were carefully examined for the presence of functional domains (Pfam and Interpro databases) that could support their involvement in cholesterol transport, conversion, or metabolism.

M. leprae putative ChoD (ML0389) and putative 3 β -hydroxysteroid dehydrogenase (3 β -HSD) (ML1942), *M. tuberculosis* putative ChoD (Rv3409c) and 3 β -HSD (Rv1106c), and *Streptomyces* sp. 3 β -HSD (Pro-

tein Data Bank [PDB] entry 1B4V) sequences were analyzed using bioinformatics tools for protein sequence annotation. Sequence similarity searches in the nonredundant and PDB databases were performed using online NCBI BLASTP tool. Multiple alignments were performed using Clustal Omega (<http://www.ebi.ac.uk>). The NCBI conserved-domain database (CDD), PFAM (<http://pfam.xfam.org>), and Interpro (<http://www.ebi.ac.uk/interpro/>) were searched for the identification of functional domains and active site residues. Subcellular localization of proteins was predicted with PSORTb (<http://www.psорт.org/psортb/>) and MycoSub (27). Secondary structures were predicted using PredictProtein (<https://www.predictprotein.org>).

Statistical analysis. An unpaired *t* test was performed using the GraphPad InStat program (GraphPad Software, San Diego, CA) and *P* values of <0.05 were considered statistically significant.

RESULTS AND DISCUSSION

In silico analysis of *M. leprae* putative cholesterol metabolism genes. *M. leprae* possesses a genome with less than one-third of the functional genes annotated in the *M. tuberculosis* genome (28), thus retaining what is considered the smallest set of genes necessary for mycobacterial pathogenicity. The annotated *M. tuberculosis* genome contains approximately 250 genes involved in lipid and fatty acid metabolism, and transcriptional profiling of *M. tuberculosis* cultured with and without cholesterol identified over 200 genes regulated in response to cholesterol (20). Since intracellular *M. leprae* survival depends on cholesterol (12), we performed an *in silico* search for *M. leprae* orthologs of the *M. tuberculosis* enzymes/proteins associated with cholesterol metabolism. From the 139 *M. tuberculosis* gene products associated with cholesterol metabolism (21), only 11 were encoded in the *M. leprae* genome (Table 1). Out of this list, 3 β -HSD was likely the only gene product dedicated exclusively to steroid metabolism. The 3 β -HSD enzyme of *M. tuberculosis* has been biochemically validated and shown to oxidize cholesterol to cholestenone, the first step of ring degradation (29) (Fig. 1A). Most of the remaining 10 *M. leprae* gene products were potential enzymes involved in the beta oxidation cycle, predicted to participate in the degradation of both the cholesterol side chain and fatty acids (21).

The remaining set of enzymes involved in the sterol ring and side chain degradation located in the “*cho* island” of *M. tuberculosis* were not found in *M. leprae* genome or were identified as pseudogenes. Also, the *mce4* operon, which codes for an active transport system for sterol lipids in mycobacteria, including cholesterol (30), is absent in *M. leprae*. Taken together, the results of this *in silico* analysis suggest that *M. leprae* is able to oxidize cholesterol to cholestenone but unable to further degrade its sterol rings and side chain.

***M. leprae* is unable to utilize cholesterol as a nutrition source.** The capacity of *M. leprae* to metabolize cholesterol was further investigated by incubating suspensions of freshly isolated athymic *nu/nu* mouse-derived *M. leprae* with radiolabeled cholesterol as the sole carbon source. Both [4-¹⁴C]cholesterol and [26-¹⁴C]cholesterol, respectively, were used to access the capacity of *M. leprae* to degrade the sterol ring and the side chain of cholesterol. As a positive control, *M. leprae* was incubated with [1-¹⁴C]palmitic acid, the carbon source used in radiorespirometry assays to measure bacterial viability (31). Live *M. tuberculosis* also served as a positive control, and heat-inactivated *M. leprae* or *M. tuberculosis* was used as a negative control. *M. leprae* cells were incubated for 4 days with the above-mentioned radiolabeled substrates, and the amount of radioactivity incorporated into the bac-

TABLE 1 *Mycobacterium leprae* genes exhibiting high similarity to *M. tuberculosis* genes involved in cholesterol metabolism

<i>M. leprae</i> locus	Gene	Protein description ^d	<i>M. tuberculosis</i> locus	% similarity	% coverage
ML0348	<i>ml0348</i>	Possible coenzyme F ₄₂₀ -dependent oxidoreductase	Rv3520c ^a	87	97
ML0354	<i>ilvX</i>	Putative acetoacetyl-CoA synthase I large subunit	Rv3509c ^b	87	100
ML0559	<i>ribA</i>	Putative GTP cyclohydrolase II/3,4-dihydroxy-2-butanone-4-phosphate synthase	Rv1940b	96	97
ML0660	<i>fadE23</i>	Putative acyl-CoA dehydrogenase	Rv3140a	94	100
ML0661	<i>fadE24</i>	Putative acyl-CoA dehydrogenase	Rv3139a	90	99
ML1051	<i>xclC</i> (<i>fadD36</i>) ^e	Acyl-CoA synthase	Rv1193b	88	100
ML1158	<i>fadA4</i>	Possible acetyl-CoA C-acetyltransferase	Rv1323b	93	98
ML1942	<i>hsd</i>	Probable cholesterol dehydrogenase	Rv1106c	85	99
ML2401	<i>echA9</i>	Putative enoyl-CoA hydratase/isomerase	Rv1071c ^a	85	100
ML2461	<i>fadB2</i>	3-Hydroxyacyl-CoA dehydrogenase	Rv0468c	94	100
ML2563	<i>fadE5</i>	Acyl-CoA dehydrogenase	Rv0244c ^b	94	100

^a Biochemically predicted.^b Computationally annotated.^c Biochemically validated.^d CoA, coenzyme A.^e *xclC* is alternatively designated *fadD36*.

terial cells was measured. As shown in Fig. 1B, both palmitic acid and cholesterol were incorporated by *M. leprae*. Only live, not heat-inactivated, bacteria were able to accumulate palmitic acid. In contrast, heat-inactivated *M. leprae* incorporated cholesterol to a higher level than the live cells. Exogenous cholesterol incorporation into *M. leprae* was observed as early as 30 min and reached a maximum level at 24 h of incubation. Identical incorporation kinetics were observed with live and heat-inactivated bacteria (data not shown). Since *M. leprae* has lost the operon (*mce4*) encoding the active transport system for sterols, the observed cholesterol incorporation probably occurs at the cell envelope level

through interaction of cholesterol with cell wall components. An explanation for the higher levels of cholesterol incorporation by heat-inactivated *M. leprae* could be alteration of the cell wall architecture by heat, resulting in the exposure of sites with high-affinity binding of cholesterol. A potential binding site for cholesterol in *M. leprae* could be the mycolic acids, a major constituent of the mycobacterial cell wall. According to Benadie et al. (32), the mycolic acids mimic the structure of cholesterol and exhibit an affinity for it, being able to interact with cholesterol. These observations could explain the cholesterol incorporation even by heat-inactivated bacteria.




FIG 1 Cholesterol is incorporated into *M. leprae* cells but is not used as an energy or carbon source. (A) Chemical structures and reaction of the oxidation of cholesterol to cholestenone catalyzed by 3 β -HSD in *M. tuberculosis*. Asterisks indicate the position of the radiolabel (carbon-4 and carbon-26) in the molecules. The chemical structures were designed using Chem Draw (PerkinElmer). (B to D) Heat-inactivated (HI) and live (L) *M. leprae* (6×10^7 bacteria/200 μ l) were incubated for 4 days at 33°C with 1 μ Ci/ml [4-¹⁴C]cholesterol, [26-¹⁴C]cholesterol, or [1-¹⁴C]palmitic acid (PA). (B) Bacteria were washed, and incorporation of radiolabeled lipids into *M. leprae* cells was expressed as kilodisintegrations per minute (kDPM) per milligram of protein. (C) For the radiorespirometry assay, the ¹⁴CO₂ trapped on strips was measured and expressed as DPM. Data are means and standard deviations of the results of three experiments performed in duplicate. (D) TLC of lipid extracts of *M. leprae* cells labeled with [4-¹⁴C]cholesterol, [26-¹⁴C]cholesterol, or [1-¹⁴C]palmitic acid. Labeled *M. leprae* cells were washed, extracted with chloroform-methanol (2:1), and resolved by TLC analysis with chloroform-methanol (95:5). Radiolabeled lipids were detected using a PhosphorImager. All data are representative of three independent experiments performed in duplicate. SF, solvent front; Chon, cholestenone; Cho, cholesterol; PA, palmitic acid; PGL-I, phenolic glycolipid I. **, $P = 0.0013$, and *, $P < 0.05$, for heat-inactivated versus live *M. leprae*.




FIG 2 *M. leprae* oxidizes cholesterol to cholestenone. (A) Comigration on TLC of radiolabeled compounds extracted from *M. leprae* with cholesterol and cholestenone (a and c). Cholesterol and cholestenone standards (2 μ g of each) were used to spike *M. leprae* lipid extracts and resolved by TLC with solvent I (chloroform-methanol-acetic acid [95:4:1]) or solvent II (chloroform-methanol-ammonium hydroxide [95:4:1]). Cold standards were visualized by spraying the same plates with a 5% phosphomolybdic acid spray and heating them with a heat gun until the bands were visible (b and d). HI, heat inactivated; L, live. (B) Evidence of cholestenone in the spent medium. Bacilli were incubated with [4- 14 C]cholesterol for different time periods at 33°C and separated from spent medium by centrifugation followed by extensive washing with PBS-tyloxapol. Washed bacilli (lanes B) as well as spent medium (lanes M) were extracted with chloroform-methanol (2:1), and the resulting lipids were analyzed by TLC using chloroform-ammonium acetate (97:3) as the running solvent. All TLC plates were exposed to the phosphor screen and radiolabeled compounds were visualized by PhosphorImager analysis. Standards were visualized by spraying the TLC plates with anisaldehyde solution and heating. SF, solvent front; Chon, cholestenone; Cho, cholesterol; AD, androstenedione. (C) Time course of *M. leprae* cholestenone production and secretion. *M. leprae* was incubated with [4- 14 C]cholesterol for 0, 6, 12, 24, and 48 h at 33°C. Total lipid extraction and TLC analysis from bacteria and spent medium were performed as described. Densitometry analysis of the chromatogram was performed using the Image QuantTL program. Data are representative of two independent experiments.

Next, the capacity of *M. leprae* to oxidize [4- 14 C]cholesterol or [26- 14 C]cholesterol, generating 14 CO₂, was measured by a radiorespirometry assay. Moreover, potential utilization of cholesterol for the biosynthesis of bacterial lipids was assessed by extraction of *M. leprae* lipids and analysis by TLC. As shown in Fig. 1C, live but not heat-inactivated *M. leprae* oxidized [1- 14 C]palmitic acid, producing 14 CO₂. In contrast, increased levels of 14 CO₂ production above the background were not observed in the presence of [4- 14 C]cholesterol or [26- 14 C]cholesterol (Fig. 1C). Palmitic acid-derived carbon was also efficiently incorporated into *M. leprae* lipids, including the species-specific PGL-I, after 4 days of incubation (Fig. 1D). However, when incubated with [4- 14 C]cholesterol or [26- 14 C]cholesterol, *M. leprae* catalyzed the formation of only one 14 C-labeled TLC spot in addition to cholesterol. This product migrated with a retardation factor (Rf) consistent with that of

cholestenone. Parallel assays were conducted with *M. tuberculosis* as a control using identical protocols, but with shorter periods of incubation (24 and 72 h). As reported previously, *M. tuberculosis* was able to utilize cholesterol (18) as well as palmitic acid as a source of energy and carbon (see Fig. S1 in the supplemental material). Altogether, the results generated confirmed the *in silico* prediction that *M. leprae* is unable to utilize both the sterol rings and the side chain of cholesterol as nutritional sources.

***M. leprae* oxidizes cholesterol to cholestenone.** To confirm the identity of the product generated from exogenous cholesterol by *M. leprae*, unlabeled cholesterol and cholestenone standards were spiked into live and heat-inactivated *M. leprae* radiolabeled lipid extracts. The spiked samples were analyzed by TLC using both acidic (Fig. 2A, solvent I) and basic (Fig. 2A, solvent II) solvent conditions. As shown in Fig. 2A and Table 2, the Rf of the

TABLE 2 Retardation factor (Rf) values of the radiolabeled compound produced by *M. leprae* incubated with [4-¹⁴C]cholesterol

Substance	Rf value for solvent ^a							
	I				II			
	Standard		Labeled sample		Standard		Labeled sample	
	HI	L	HI	L	HI	L	HI	L
Cholesterol	0.52	0.51	0.52	0.51	0.72	0.72	0.74	0.74
Cholestenone	0.77	0.78	0.78	0.78	0.90	0.90	0.91	0.91

^a Solvent I, chloroform-methanol-acetic acid (95:4:1); solvent II, chloroform-methanol-ammonium hydroxide (95:4:1); L, live; HI, heat inactivated.

lipid produced by *M. leprae* from [4-¹⁴C]cholesterol corresponded to the cholestenone standard. Interestingly, labeled cholestenone was readily extracted from spent medium as well as cells at 2 to 7 days of incubation, as shown in Fig. 2B. Androstenedione (AD), the steroid intermediate generated after the complete degradation of the cholesterol side chain, was not detected in any sample. In concordance with the rapid cholesterol incorporation into the bacterial cells, cholestenone generation was observed as early as 6 h after incubation with cholesterol (Fig. 2C). This rapid conversion of cholesterol to cholestenone, in the likely absence of a cholesterol transport mechanism, suggests that cholestenone formation occurs at the bacterial cell surface.

Prediction of the *M. leprae* cholesterol oxidase. The first reaction of ring metabolism is the oxidation and isomerization of cholesterol to form cholestenone (Fig. 1A). In bacteria, this process is catalyzed by either 3 β -HSD or cholesterol oxidase. Al-

though these enzymes utilize distinct reaction mechanisms, they catalyze the same transformation, and both can be found within the genomes of steroid-utilizing bacteria (21). The capacity of *M. leprae* to oxidize cholesterol to cholestenone was expected based on the annotation of an intact *hsd* gene (*ml1942*). The *M. leprae* 3 β -HSD ortholog is 75% identical and 85% similar to the *M. tuberculosis* 3 β -HSD (Rv1106c) (Table 1), whose cholesterol oxidation activity was experimentally confirmed (29). ML1942 contains a conserved NAD binding motif, as well as a conserved active site and substrate binding residues (Fig. 3). The nonconserved substitutions between *M. leprae* and *M. tuberculosis* proteins do not result in significant changes in the predicted secondary structure (data not shown). Thus, the bioinformatics of ML1942 provides evidence that this protein functions as a 3 β -HSD that would allow *M. leprae* to convert cholesterol to cholestenone.

A previous proteomics study showed the presence of the enzyme associated with the cell wall of *M. leprae* (33). This subcellular localization is consistent with the current data that demonstrated the absence of the cholesterol import machinery in *M. leprae* and rapid accumulation of cholestenone in the spent culture medium, supporting the hypothesis that cholesterol transformation to cholestenone occurs at the bacterial surface.

Besides the *hsd* gene, the genomes of *M. leprae* and *M. tuberculosis* encode annotated ChoD cholesterol oxidase proteins (ML0389 and Rv3409c, respectively). The *M. leprae* ChoD possesses high sequence similarity to the *M. tuberculosis* ChoD (see Fig. S2 in the supplemental material). However, alignment of these two sequences to a functional ChoD (PDB ID 1B4V) from *Streptomyces* sp. revealed an absence of amino acid residues composing the well-defined FAD binding and active sites required for

```

ML  MLRQVGNASLTTELGRVLVTGGSGFVGANLVTALLERGYQVRSFDRAPMPLPQHPQLEVL
TB  MLRMGDASLTTELGRVLVTGGAGFVGANLVTLLDRGHWVRSFDRAPSLLPAHPQLEVL
    ***  :*:*****:*****:***:***: *****  *  *****

ML  QGDITDATVCTTAMDSIDTVFHTAAIIELMGGASVTDEYRQRSYTVVGGTENLLRAGQK
TB  QGDITDADVCAAAVDGIDTIFHTAAIIELMGGASVTDEYRQRSFAVVGGTENLLHAGQR
    *****  *  :*: * . ***:*****:*****:*****:*****:*****:***:

ML  SGVKRFVYTASNSVVMGGTPITGGDETMPYTKRFNDLYTETHVVAEKFVLSQNGVPDGET
TB  AGVQRFVYTSSNSVVMGGQNIAGGDETLPYTDRFNDLYTETHVVAERFVLAQNGVD---G
    :*:*****:*****  *  :*****:***:*****:*****:*****:*****:

ML  MLTCSIRPSGIWGRGDQTMRKAFESVSVSGHVKVLIGSKNAKLDNSYVHNLVHGLILAAE
TB  MLTCAIRPSGIWNGNDQTMRKLFESVLKGHVKVLVGRKSARLDNSYVHNLIHGFILAAA
    *****:*****.*****  ***: . *****: *  * .:*****:***:*****

ML  HLVPGGTAPGQAYFINDGEPINFFFMGPIIKACGENWPRVRISGRLVRNVMAVWQRLHF
TB  HLVPDGTAPGQAYFINDAEPINMFEFARPVLEACGQRWPKMRISGPAVRVMTGWQRLHF
    ****  *****:*****.*****:***  *  :*:***: . ***:*****  **  *  :  *****

ML  GFGLPKPMEPLAVERVYLDNYFSIEKAHKELGYRPLFTTEQAMAECLPYTELFEIQKV
TB  RFGFPALLEPLAVERLYLDNYFSIAKARRDLGEYEPLFTTQQALTECLPYVSLFEQMKN
    ***  *  *  :*****:*****  ***: .:***.*****:***:*****. . *****  *

ML  AAKPHLASIAATPRE
TB  EARAEKTAATVKP---
    *  .  .  :  :  .  .  *

```

FIG 3 3 β -Hydroxysteroid dehydrogenase protein sequences from *M. leprae* (ML1942) and *M. tuberculosis* (Rv1106c). Asterisks indicate positions which have identical amino acid residues, colons indicate conserved amino acid substitution between groups with strongly similar properties, and periods indicate amino acid conservation between groups with weakly similar amino acid properties, according to Clustal software. Yellow highlighting indicates the NAD binding motif, green highlighting indicates the active-site residues, and blue highlighting indicates additional substrate binding residues. Nonconserved substitutions are indicated by colored amino acid residues, according to the Clustal color code: red, small and hydrophobic; blue, acidic; magenta, basic; green, containing hydroxyl, sulphydryl, and amine. ML, *M. leprae*; TB, *M. tuberculosis*.




FIG 4 Cholestenone is produced *in vivo* by *M. leprae*. (A) GC-MS chromatogram of bis(trimethylsilyl)trifluoroacetamide (BSTFA)-trimethylchlorosilane (TMCS)-derived extracts of *M. leprae*; (B) steroid standards mix. *M. leprae* was isolated from footpads of infected mice and submitted to total lipids extracted with chloroform-methanol (2:1). Dried lipids were saponified with 25% alcoholic potassium hydroxide solution, extracted with heptane, and derivatized with BSTFA-TMCS (99:1) and pyridine. Standards: 1, cholesterol; 2, cholesta-5,7-dien-3-ol; 3, cholesta-8,24-dien-3-ol; 4, ergosta-5,7,22-trien-3-ol; 5, cholest-4-en-3-one; 6, stigmasterol; 7, lanosta-8,24-dien-3-ol; 8, β -sitosterol.

cholesterol oxidase activity (see Fig. S2 in the supplemental material). In addition, the two loop regions of *Streptomyces* sp. and *Brevibacterium* sp. cholesterol oxidases involved in positioning the substrate in the active site are weakly conserved in putative mycobacterial ChoD proteins. Furthermore, ChoD shows a low level of amino acid identity ($\sim 24\%$) with cholesterol oxidases from other bacteria as well as other glucose-methanol-choline (GMC) oxidoreductase superfamily members; only the FAD binding region is conserved (34, 35). Moreover, a recent study by Gao and Sampson (36) suggests that Rv3409c controls acetylation of cell surface glycopeptidolipids, which affects activation of the innate immune system. Experimental evidence also indicates that the *M. tuberculosis* ChoD does not catalyze the transformation of cholesterol to cholestenone (37); in addition, an *M. tuberculosis* strain with a deletion of the *hsd* gene was incapable of oxidizing cholesterol (29, 37). This provides further evidence that 3β -HSD is the sole cholesterol-oxidizing enzyme in *M. tuberculosis* and that the ChoD of *M. leprae* and *M. tuberculosis* may have a function other than cholesterol oxidation.

***In vivo* production of cholestenone by *M. leprae*.** The *M. tuberculosis* 3β -HSD recombinant enzyme was shown to also oxidize other 3-hydroxysterols, such as pregnenolone and dehydroepiandrosterone, to their respective 3-keto-4-ene products (29). Thus, we investigated whether the *in vitro* transformation of cholesterol to cholestenone by *M. leprae* occurs during *in vivo* infection. *M. leprae* isolated from footpads of infected mice was subjected to lipid extraction followed by GC-MS analysis. Two peaks

of cholesterol were detected, the first corresponding to nonsilylated cholesterol (Fig. 4A). This observation correlated with the *in vitro* data that demonstrated the accumulation of cholesterol on the cell surface of *M. leprae*. A smaller peak with a retention time and mass identical (Fig. 4A; also, see Fig. S3 in the supplemental material) to those of cholestenone was also detected in the footpad-derived *M. leprae*. Since trace amounts of cholestenone can be found in mammalian tissue (38), the possibility that the detected cholestenone was derived from contaminating host tissue was assessed by analyzing lipids directly extracted from uninfected mouse footpad tissue. As shown in Fig. S4A in the supplemental material, no cholestenone was detected in uninfected tissue, indicating that the cholestenone found in *M. leprae* purified from the mouse footpad was truly a product of *in vivo* bacterial metabolism.

The retention of the *hsd* gene in the genome of *M. leprae*, despite the reductive evolution of *M. leprae*, and the observed production of cholestenone by *M. leprae* in infected tissue suggests a role for sterol-oxidizing activity in bacterial pathogenesis. We recently revealed that *M. leprae* infection induces the expression and activation of sterol regulatory element-binding protein (SREBP), a transcription factor that regulates the biosynthesis and uptake of cholesterol (12, 39). Liver X receptor (LXR), another important transcription factor, provides a regulatory system for the elimination of excess cholesterol (40). The activities of the SREBP and LXR pathways are regulated by several sterols and oxysterols that also influence innate and adaptive immune responses in chronic diseases (41). One of the most

abundant oxysterols is 27-hydroxycholesterol (27HC), which is generated by the enzyme sterol 27-hydroxylase (CYP27A1) (42, 43). *In vitro* studies have shown that cholestenone is hydroxylated at a much higher rate than corresponding sterols with a 3-hydroxy- Δ^5 structure (44). Thus, there is a possibility that the host cell CYP27A1 can oxidize cholestenone produced by *M. leprae*. Furthermore, recent data have suggested that an excess of cholestenone affects host cell membrane functionality (45–48). Therefore, we propose that cholestenone production by *M. leprae* might modulate host cell functions to facilitate *M. leprae* invasion and persistence in those cells.

Finally, the accumulation of cholesterol *per se* can play an important role in bacterial pathogenesis. Recently, different roles for cholesterol during microbial infection as well as in cell processes have been described. Cholesterol increases *Helicobacter pylori* resistance to several antibiotics, such as tetracycline and clarithromycin, due to its incorporation in the bacterial membrane (49). It has been reported that *M. tuberculosis* incorporates cholesterol into the cell wall, leading to decreased rifampin uptake (19). This phenomenon could explain the enhanced mycobacterial killing during the combined treatment of *M. leprae*- and *M. tuberculosis*-infected macrophages with atorvastatin and rifampin (15). Furthermore, as mentioned before, cholesterol may interact with mycolic acids in the cell wall (32), leading to an even more decreased cell wall permeability, since cholesterol has major implications in membrane fluidity and rigidity. Considering that rifampin is able to cross the cell wall due to its hydrophobicity (50), any alterations in the cell wall permeability can directly impact rifampin on uptake, as observed by Brzostek et al. (19). In addition, cholesterol accumulation has been implicated in inhibiting phagosome fusion with lysosomes (51), and reports showed that *M. leprae* inhibits phagolysosome fusion, which contributes to its survival inside the host cell (52). Taken together, this evidence supports the idea that cholesterol accumulation influences, in multiple ways, host-pathogen interactions that might contribute to infection persistence.

In conclusion, our data indicate that *M. leprae* lost the capacity to catabolize the sterol rings and the side chain of cholesterol, preserving only the capacity to oxidize it to cholestenone, the first step of cholesterol degradation. It was also shown that cholesterol is likely incorporated into the cell envelope by a passive mechanism, since *M. leprae* has lost the *mce4* operon responsible for coding the active transport system in mycobacteria devoted to sterols. Interestingly, *M. tuberculosis* strains with mutations in *mce4* also retain some ability to incorporate cholesterol. This residual cholesterol uptake was suggested to be due to another, less efficient import system or to a passive diffusion into the cell (18). Transformation of cholesterol to cholestenone was observed to occur both *in vitro* and *in vivo* and most likely occurs at the bacterial cell surface catalyzed by the enzyme 3β -HSD. Our study underscores the complexity of sterol metabolism and that its importance for mycobacterial pathogenesis extends well beyond its utilization as a source of nutrition. This is also evident in the context of *M. tuberculosis* infection, where cholesterol accumulation in the host cell but not its degradation seems to be important for bacterial survival during the active phase of infection when multiple other carbon sources are available for sustaining bacterial intracellular growth (21, 53, 54). Thus, a future challenge is to understand the biochemical basis of the role of cholesterol,

including its oxidation step, in *M. leprae* pathogenesis. Further elucidation of cholesterol metabolism in the host cell during *M. leprae* infection will establish the mechanism by which this lipid supports *M. leprae* intracellular survival and will open new avenues for novel leprosy therapies.

ACKNOWLEDGMENTS

This work was supported by the Heiser Program for Research in Leprosy and the Brazilian Research Council (CNPq). Carlos Adriano M. Silva is a postdoctorate fellow from Science without Borders—Capes/Brazil.

We are grateful to Ramanuj Lahiri and Patricia S. Rosa for providing *M. leprae*. We thank Mileane de Souza Busch and Amanda Ward for technical support.

REFERENCES

- Han XY, Silva FJ. 2014. On the age of leprosy. *PLoS Negl Trop Dis* 8:e2544. <http://dx.doi.org/10.1371/journal.pntd.0002544>.
- WHO. 2013. Global leprosy: update on the 2012 situation. *Wkly Epidemiol Rec* 88:365–379.
- Barreto JG, Bisanzio D, Guimarães LS, Spencer JS, Vazquez-Prokopec GM, Kitron U, Salgado CG. 2014. Spatial analysis spotlighting early childhood leprosy transmission in a hyperendemic municipality of the Brazilian Amazon region. *PLoS Negl Trop Dis* 8:e2665. <http://dx.doi.org/10.1371/journal.pntd.0002665>.
- Nery JADC, Filho FB, Quintanilha J, Machado AM, Oliveira SDSC, Sales AM. 2013. Understanding the type 1 reactional state for early diagnosis and treatment: a way to avoid disability in leprosy. *An Bras Dermatol* 88:787–792. <http://dx.doi.org/10.1590/abd1806-4841.20132004>.
- Cole ST, Eglmeier K, Parkhill J, James KD, Thomson NR, Wheeler PR, Honoré N, Garnier T, Churcher C, Harris D, Mungall K, Basham D, Brown D, Chillingworth T, Connor R, Davies RM, Devlin K, Duthoy S, Feltwell T, Fraser A, Hamlin N, Holroyd S, Hornsby T, Jagels K, Lacroix C, Maclean J, Moule S, Murphy L, Oliver K, Quail MA, Rajandream MA, Rutherford KM, Rutter S, Seeger K, Simon S, Simmonds M, Skelton J, Squares R, Squares S, Stevens K, Taylor K, Whitehead S, Woodward JR, Barrell BG. 2001. Massive gene decay in the leprosy bacillus. *Nature* 409:1007–1011. <http://dx.doi.org/10.1038/35059006>.
- Virchow R. 1863. Die krankhaften Geschwülste, p 494–531. August Hirschwald, Berlin, Germany.
- Kaplan G, Van Voorhis WC, Sarno EN, Nogueira N, Cohn ZA. 1983. The cutaneous infiltrates of leprosy. A transmission electron microscopy study. *J Exp Med* 158:1145–1159.
- Brennan PJ. 1984. *Mycobacterium leprae*—the outer lipoidal surface. *J Biosci* 6:685–689. <http://dx.doi.org/10.1007/BF02702710>.
- Mattos KA, D'Avila H, Rodrigues LS, Oliveira VGC, Sarno EN, Atella GC, Pereira GM, Bozza PT, Pessolani MCV. 2010. Lipid droplet formation in leprosy: Toll-like receptor-regulated organelles involved in eicosanoid formation and *Mycobacterium leprae* pathogenesis. *J Leukoc Biol* 87:371–384. <http://dx.doi.org/10.1189/jlb.0609433>.
- Cruz D, Watson AD, Miller CS, Montoya D, Ochoa MT, Sieling PA, Gutierrez MA, Navab M, Reddy ST, Witztum JL, Fogelman AM, Rea TH, Eisenberg D, Berliner J, Modlin RL. 2008. Host-derived oxidized phospholipids and HDL regulate innate immunity in human leprosy. *J Clin Invest* 118:2917–2928. <http://dx.doi.org/10.1172/JCI34189>.
- Tanigawa K, Suzuki K, Nakamura K, Akama T, Kawashima A, Wu H, Hayashi M, Takahashi SI, Ikuyama S, Ito T, Ishii N. 2008. Expression of adipose differentiation-related protein (ADRP) and perilipin in macrophages infected with *Mycobacterium leprae*. *FEMS Microbiol Lett* 289:72–79. <http://dx.doi.org/10.1111/j.1574-6968.2008.01369.x>.
- Mattos KA, Oliveira VGC, Berrêdo-Pinho M, Amaral JJ, Antunes LCM, Melo RCN, Acosta CCD, Moura DF, Olmo R, Han J, Rosa PS, Almeida PE, Finlay BB, Borchers CH, Sarno EN, Bozza PT, Atella GC, Pessolani MCV. 2014. *Mycobacterium leprae* intracellular survival relies on cholesterol accumulation in infected macrophages: a potential target for new drugs for leprosy treatment. *Cell Microbiol* 16:797–815. <http://dx.doi.org/10.1111/cmi.12279>.
- Mattos KA, Oliveira VGC, D'Avila H, Rodrigues LS, Pinheiro RO, Sarno EN, Pessolani MCV, Bozza PT. 2011. TLR6-driven lipid droplets in *Mycobacterium leprae*-infected Schwann cells: immunoinflammatory

- platforms associated with bacterial persistence. *J Immunol* 187:2548–2558. <http://dx.doi.org/10.4049/jimmunol.1101344>.
14. Mattos KA, Lara FA, Oliveira VGC, Rodrigues LS, D'Avila H, Melo RCN, Manso PPA, Sarno EN, Bozza PT, Pessolani MCV. 2011. Modulation of lipid droplets by *Mycobacterium leprae* in Schwann cells: a putative mechanism for host lipid acquisition and bacterial survival in phagosomes. *Cell Microbiol* 13:259–273. <http://dx.doi.org/10.1111/j.1462-5822.2010.01533.x>.
 15. Lobato LS, Rosa PS, Ferreira JDS, Neumann ADS, Da Silva MG, Do Nascimento DC, Soares CT, Pedrini SCB, De Oliveira DSL, Monteiro CP, Pereira GMB, Ribeiro-Alves M, Hacker MA, Moraes MO, Pessolani MCV, Duarte RS, Lara FA. 2014. Statins increase rifampin mycobactericidal effect. *Antimicrob Agents Chemother* 58:5766–5774. <http://dx.doi.org/10.1128/AAC.01826-13>.
 16. Peyron P, Vaubourgeix J, Poquet Y, Levillain F, Botanch C, Bardou F, Daffé M, Emile JF, Marchou B, Cardona P-J, De Chastellier C, Altare F. 2008. Foamy macrophages from tuberculosis patients' granulomas constitute a nutrient-rich reservoir for *M. tuberculosis* persistence. *PLoS Pathog* 4:e1000204. <http://dx.doi.org/10.1371/journal.ppat.1000204>.
 17. Russell DG, Cardona P-J, Kim M-J, Allain S, Altare F. 2009. Foamy macrophages and the progression of the human tuberculosis granuloma. *Nat Immunol* 10:943–948. <http://dx.doi.org/10.1038/ni.1781>.
 18. Pandey AK, Sasseti CM. 2008. Mycobacterial persistence requires the utilization of host cholesterol. *Proc Natl Acad Sci U S A* 105:4376–4380. <http://dx.doi.org/10.1073/pnas.0711159105>.
 19. Brzostek A, Pawelczyk J, Rumijowska-Galewicz A, Dziadek B, Dziadek J. 2009. *Mycobacterium tuberculosis* is able to accumulate and utilize cholesterol. *J Bacteriol* 191:6584–6591. <http://dx.doi.org/10.1128/JB.00488-09>.
 20. Nesbitt NM, Yang X, Fontán P, Kolesnikova I, Smith I, Sampson NS, Dubnau E. 2010. A thiolase of *Mycobacterium tuberculosis* is required for virulence and production of androstenedione and androstadienedione from cholesterol. *Infect Immun* 78:275–282. <http://dx.doi.org/10.1128/IAI.00893-09>.
 21. Wiperman MF, Sampson NS, Thomas ST. 2014. Pathogen roid rage: cholesterol utilization by *Mycobacterium tuberculosis*. *Crit Rev Biochem Mol Biol* 49:269–293. <http://dx.doi.org/10.3109/10409238.2014.895700>.
 22. Sambandamurthy VK, Wang X, Chen B, Russell RG, Derrick S, Collins FM, Morris SL, Jacobs WR, Jr. 2002. A pantothenate auxotroph of *Mycobacterium tuberculosis* is highly attenuated and protects mice against tuberculosis. *Nat Med* 8:1171–1174. <http://dx.doi.org/10.1038/nm765>.
 23. Buddemeyer EU. 1974. Liquid scintillation vial for cumulative and continuous radiometric measurement of in vitro metabolism. *Appl Microbiol* 28:177–180.
 24. Bligh EG, Dyer WJ. 1959. A rapid method of total lipid extraction and purification. *Can J Biochem Physiol* 37:911–917. <http://dx.doi.org/10.1139/o59-099>.
 25. Folch J, Lees M, Slone Stanley GH. 1957. A simple method for the isolation and purification of total lipides from animal tissues. *J Biol Chem* 226:497–509.
 26. Dunphy PJ, Kerr JD, Pennock JF, Whittle KJ, Feeney J. 1967. The plurality of long chain isoprenoid alcohols (polyprenols) from natural sources. *Biochim Biophys Acta* 136:136–147. [http://dx.doi.org/10.1016/0304-4165\(67\)90329-7](http://dx.doi.org/10.1016/0304-4165(67)90329-7).
 27. Zhu PP, Li WC, Zhong ZJ, Deng EZ, Ding H, Chen W, Lin H. 2015. Predicting the subcellular localization of mycobacterial proteins by incorporating the optimal tripeptides into the general form of pseudo amino acid composition. *Mol Biosyst* 11:558–563. <http://dx.doi.org/10.1039/C4MB00645C>.
 28. Cole ST, Brosch R, Parkhill J, Garnier T, Churcher C, Harris D, Gordon SV, Eiglmeier K, Gas S, Barry CE, Tekaiia F, Badcock K, Basham D, Brown D, Chillingworth T, Connor R, Davies R, Devlin K, Feltwell T, Gentles S, Hamlin N, Holroyd S, Hornsby T, Jagels K, Krogh A, McLean J, Moule S, Murphy L, Oliver K, Osborne J, Quail MA, Rajandream MA, Rogers J, Rutter S, Seeger K, Skelton J, Squares R, Squares S, Sulston JE, Taylor K, Whitehead S, Barrell BG. 1998. Deciphering the biology of *Mycobacterium tuberculosis* from the complete genome sequence. *Nature* 393:537–544. <http://dx.doi.org/10.1038/31159>.
 29. Yang X, Dubnau E, Smith I, Sampson NS. 2007. Rv1106c from *Mycobacterium tuberculosis* is a β -hydroxysteroid dehydrogenase. *Biochemistry* 46:9058–9067. <http://dx.doi.org/10.1021/bi700688x>.
 30. Mohn WW, Van Der Geize R, Stewart GR, Okamoto S, Liu J, Dijkhuizen L, Eltis LD. 2008. The actinobacterial *mce4* locus encodes a steroid transporter. *J Biol Chem* 283:35368–35374. <http://dx.doi.org/10.1074/jbc.M805496200>.
 31. Franzblau SG. 1988. Oxidation of palmitic acid by *Mycobacterium leprae* in an axenic medium. *J Clin Microbiol* 26:18–21.
 32. Benadie Y, Deysel M, Siko DGR, Roberts VV, Van Wyngaardt S, Thanyani ST, Sekanka G, Ten Bokum AMC, Collett LA, Grooten J, Baird MS, Verschoor JA. 2008. Cholesteroid nature of free mycolic acids from *M. tuberculosis*. *Chem Phys Lipids* 152:95–103. <http://dx.doi.org/10.1016/j.chemphyslip.2008.01.004>.
 33. Marques MAM, Neves-Ferreira AGC, Xavier Da Silveira EK, Valente RH, Chapeaurouge A, Perales J, Bernardes RDS, Dobos KM, Spencer JS, Brennan PJ, Pessolani MCV. 2008. Deciphering the proteomic profile of *Mycobacterium leprae* cell envelope. *Proteomics* 8:2477–2491. <http://dx.doi.org/10.1002/pmic.200700971>.
 34. Uhia I, Galán B, Morales V, García JL. 2011. Initial step in the catabolism of cholesterol by *Mycobacterium smegmatis* mc²155. *Environ Microbiol* 13:943–959. <http://dx.doi.org/10.1111/j.1462-2920.2010.02398.x>.
 35. Cavener DR. 1992. GMC oxidoreductases: a newly defined family of homologous proteins with diverse catalytic activities. *J Mol Biol* 223:811–814. [http://dx.doi.org/10.1016/0022-2836\(92\)90992-S](http://dx.doi.org/10.1016/0022-2836(92)90992-S).
 36. Gao J, Sampson NS. 2014. A GMC oxidoreductase homologue is required for acetylation of glycopeptidolipid in *mycobacterium smegmatis*. *Biochemistry* 53:611–613. <http://dx.doi.org/10.1021/bi4015083>.
 37. Yang X, Gao J, Smith I, Dubnau E, Sampson NS. 2011. Cholesterol is not an essential source of nutrition for *Mycobacterium tuberculosis* during infection. *J Bacteriol* 193:1473–1476. <http://dx.doi.org/10.1128/JB.01210-10>.
 38. Gray MF, Lawrie TDV, Brooks CJW. 1971. Isolation and identification of cholesterol α -oxide and other minor sterols in human serum. *Lipids* 6:836–843. <http://dx.doi.org/10.1007/BF02531214>.
 39. Horton JD, Shah NA, Warrington JA, Anderson NN, Park SW, Brown MS, Goldstein JL. 2003. Combined analysis of oligonucleotide microarray data from transgenic and knockout mice identifies direct SREBP target genes. *Proc Natl Acad Sci U S A* 100:12027–12032. <http://dx.doi.org/10.1073/pnas.1534923100>.
 40. Tontonoz P, Mangelsdorf DJ. 2003. Liver X receptor signaling pathways in cardiovascular disease. *Mol Endocrinol* 17:985–993. <http://dx.doi.org/10.1210/me.2003-0061>.
 41. Spann NJ, Glass CK. 2013. Sterols and oxysterols in immune cell function. *Nat Immunol* 14:893–900. <http://dx.doi.org/10.1038/ni.2681>.
 42. Brown AJ, Jessup W. 1999. Oxysterols and atherosclerosis. *Atherosclerosis* 142:1–28. [http://dx.doi.org/10.1016/S0021-9150\(98\)00196-8](http://dx.doi.org/10.1016/S0021-9150(98)00196-8).
 43. Russell DW. 2000. Oxysterol biosynthetic enzymes. *Biochim Biophys Acta* 1529:126–135. [http://dx.doi.org/10.1016/S1388-1981\(00\)00142-6](http://dx.doi.org/10.1016/S1388-1981(00)00142-6).
 44. Norlin M, von Bahr S, Björkhem I, Wikvall K. 2003. On the substrate specificity of human CYP27A1: implications for bile acid and cholesterol formation. *J Lipid Res* 44:1515–1522. <http://dx.doi.org/10.1194/jlr.M300047-JLR200>.
 45. Neuvonen M, Manna M, Morkkila S, Javanainen M, Rog T, Liu Z, Bittman R, Vattulainen I, Ikonen E. 2014. Enzymatic oxidation of cholesterol: properties and functional effects of cholestenone in cell membranes. *PLoS One* 9:e103743. <http://dx.doi.org/10.1371/journal.pone.0103743>.
 46. Bittman R, Kasireddy CK, Mattjus P, Slotte JP. 1994. Interaction of cholesterol with sphingomyelin in monolayers and vesicles. *Biochemistry* 33:11776–11781. <http://dx.doi.org/10.1021/bi00205a013>.
 47. Xu X, London E. 2000. The effects of sterol structure on membrane lipid domains reveals how cholesterol can induce lipid domain formation. *Biochemistry* 39:843–849. <http://dx.doi.org/10.1021/bi992543v>.
 48. Samsonov AV, Mihalyov I, Cohen FS. 2001. Characterization of cholesterol-sphingomyelin domains and their dynamics in bilayer membranes. *Biophys J* 81:1486–1500. [http://dx.doi.org/10.1016/S0006-3495\(01\)75803-1](http://dx.doi.org/10.1016/S0006-3495(01)75803-1).
 49. McGee DJ, George AE, Trainor EA, Horton KE, Hildebrandt E, Testerman TL. 2011. Cholesterol enhances *Helicobacter pylori* resistance to antibiotics and LL-37. *Antimicrob Agents Chemother* 55:2897–2904. <http://dx.doi.org/10.1128/AAC.00016-11>.
 50. Lambert PA. 2002. Cellular impermeability and uptake of biocides and antibiotics in gram-positive bacteria and mycobacteria. *J Appl Microbiol* 92:46S–54S. <http://dx.doi.org/10.1046/j.1365-2672.92.5s1.7.x>.
 51. Huynh KK, Gershenzon E, Grinstein S. 2008. Cholesterol accumulation

- by macrophages impairs phagosome maturation. *J Biol Chem* **283**:35745–35755. <http://dx.doi.org/10.1074/jbc.M806232200>.
52. Frehel C, Rastogi N. 1987. *Mycobacterium leprae* surface components intervene in the early phagosome-lysosome fusion inhibition event. *Infect Immun* **55**:2916–2921.
53. de Carvalho LPS, Fischer SM, Marrero J, Nathan C, Ehrt S, Rhee KY. 2010. Metabolomics of *Mycobacterium tuberculosis* reveals compartmentalized co-catabolism of carbon substrates. *Chem Biol* **17**:1122–1131. <http://dx.doi.org/10.1016/j.chembiol.2010.08.009>.
54. Beste DJV, Nöh K, Niedenführ S, Mendum TA, Hawkins ND, Ward JL, Beale MH, Wiechert W, McFadden J. 2013. ¹³C-flux spectral analysis of host-pathogen metabolism reveals a mixed diet for intracellular *mycobacterium tuberculosis*. *Chem Biol* **20**:1012–1021. <http://dx.doi.org/10.1016/j.chembiol.2013.06.012>.