

Document-ID: 02

Patron: comut

Note:

NOTICE:

Pages: 9 Printed: 01-07-11 11:15:56

Sender: Ariel/Windows

ENTEROVIRUS ISOLATION FROM FOETAL AND PLACENTAL TISSUES

N. G. S. BASSO¹, M. E. F. FONSECA¹, A. G. P. GARÇA², J. A. T. ZUARDI²,
M. R. SILVA¹, H. OUTANI²

¹Department of Virology, Institute of Microbiology, Federal University, 08040 Rio de Janeiro, and ²Fernandes Figueira Institute, Rio de Janeiro, Brazil

Received October 21, 1988

Summary. — Four spontaneous abortions and two stillbirth occurred during a prospective survey following the teratogenicity of echoviruses in 80 pregnant women selected at random from the Antenatal Care Service. Echovirus types 19, 27, and 33, Coxsackie B₂ and B₆ were isolated from placental and foetal tissues (brain, liver, kidney, heart, and spleen). The mothers also excreted the virus by faeces at least twenty days before abortion and responded serologically, indicating active virus infection. Almost all aborted children were anomalous with signs of viral infection.

Key words: enteroviruses; congenital infections; viral placentitis; foetal and placental tissues

Introduction

Since 1941 when Norman Gregg established the teratogenic role of rubella virus for humans, several authors have recognized viruses causing foetal infections (Aycok, 1941; Bates, 1955; Holowach *et al.*, 1957; Greenwald, 1958; Rhodes, 1960). Infections by cytomegalovirus, herpes, rubella, varicella-zoster, vaccinia, variola, polio, and Coxsackie B viruses during pregnancy may damage the human foetus (Hanshaw and Dudgeon, 1978).

The reports on viruses potentially causing congenital defects, stillbirth, or neonatal complications are still controversial. Of these, echoviruses have caught our attention, because their adverse effect on the foetus has not been accepted yet, although their capacity to cross the placenta had already been demonstrated (Rantsalo *et al.*, 1960; Kleinman *et al.*, 1962; Berkovich and Smithwick, 1968).

Hovata *et al.* (1983) analysing the intrauterine deaths during a period of 6 years, considering the clinical and autopsy findings, registered a case of

The work was performed at Departamento de Virologia e Setor de Microscopia Electronica de Instituto de Microbiologia da UFRJ (Brazil). Part of the Thesis was submitted by Basso, N. to the Instituto de Microbiologia, UFRJ, in partial fulfillment of the requirements for the MSc degree, supervised by Fonseca, M. E. F.

enterovirus isolation from tissues of a stillborn. In Brazil, echovirus and Coxsackie B were isolated from placentas and foetal organs (Fonseca *et al.*, 1982). There have been reports dealing with the occurrence of fatal infections "in utero", caused by echoviruses types 9 and 11 (Landsman *et al.*, 1964; Brown and Karunas, 1977; Jones *et al.*, 1980; Modlin, 1986). The present paper is the result of a prospective study made with a group of pregnant women, where the objective was the detection of maternal infection by echoviruses during the period of pregnancy and its possible damage for the foetus.

Materials and Methods

Patients. The studied population consisted of a group of 78 pregnant women that had been enrolled to the Antenatal Care Service from IFF in Rio de Janeiro, Brazil. The women were observed during the whole gestation period. Stool samples were collected monthly and two blood samples were taken: one at the beginning and one at the end of pregnancy for viral and serological studies, respectively. Placental tissue samples were collected, during delivery, for viral, histopathological, and electron microscopic studies. In stillbirth cases, tissue samples were collected from organs such as liver, brain, spleen, kidney, and lungs.

Virologic studies. Suspensions from placentas, foetal organs, and stool samples were inoculated into following cell cultures: LLC-MK₂, VERO, and HEP₂. When positive for cytopathogenic effects, cell suspensions were titrated and the viruses identified in LLC-MK₂ cells with Melnick's standard sera against enteroviruses.

Serological studies. Both blood samples collected from each mother were assayed against the virus isolated from them by neutralization test. Assays were performed to exclude other concurrent infections such as rubella, cytomegalovirus, toxoplasmosis, syphilis, listeriosis, and brucellosis. In addition, data were determined such as Rh factor, glycaemia, and blood pressure that could be important during delivery.

Immunofluorescence tests were made in paraffin-embedded tissue samples. The sections were deparaffinized with xylene (Merck) for 20 min, washed several times with PBS pH 7.2, overlaid with specific purified rabbit antiserum against the isolated viruses for one hour at 37 °C. The sections were then washed with PBS pH 7.2, covered with calf antiserum against rabbit serum labelled with fluoresceine (Boehringer Laboratories). After one hour incubation at 37 °C, the sections were washed with PBS, air dried, and mounted (Lennette and Schmidt, 1979).

Electron microscopy. The fresh tissues immediately after labour were cut into small pieces and fixed in 2.5% glutaraldehyde (Merck) for 2 hr, washed with pH 7.2 cacodylate-sucrose buffer, post-fixed by osmium tetroxide (Sigma) dehydrated in a series of alcohols and embedded in Polilyte resine (Resana). The resulting blocks were cut in an ultramicrotome. The sections were contrasted with uranyl acetate and lead citrate (Merck) and observed in electron microscope EM 301, Philips.

Results

Two stillbirth cases and three spontaneous abortions occurred in the selected group. The abortions took place between the third (one case) and fifth month of pregnancy (two cases). Five per cent (5%) of the studied pregnancies were lost by spontaneous abortion and 2.5% by stillbirth. The virus was often isolated from placenta (64.1%) rather than from faecal specimens (37.8%). About 47% of the foetal organs collected at autopsy harboured the viruses. Enterovirus had been isolated from 3 maternal faecal specimens, just before miscarriage; the same viruses were isolated from placentae and all respective foetal organs such as brain, liver, kidney, and heart. Echoviruses types 19, 27, and 33 had been isolated from 4 cases

Table 1. Abortions with enterovirus isolation

Cases	Gestational period	Occurrence	Virus isolation from	Virus type	Serology		Toxo	Syph	Brucel		
					NT	CF / CMV				ELISA/Rubella	
										1st serum	2nd serum
I	5th month	Spontaneous	Placenta	Echovirus	0	640	64	64	64		

Table 1. Abortions with enterovirus isolation

Cases	Gestational period	Occurrence	Virus isolation from	Virus type	NT		Serology		Toxo	Syph	Brucel
					1st serum	2nd serum	CF	/ CMV			
					640	64	64	64			
1	5th month	Spontaneous abortion	Placenta, faeces (10 days before abortion), brain, liver kidney	Echovirus 33	0	640	64	64	+	+	-
2	5th month	Spontaneous abortion	Placenta, faeces (a week before abortion), liver, brain	Echovirus 33	80	640	8	8	+	+	-
3	3rd month	Spontaneous abortion	Placenta, liver, brain	Coxsackie B ₂	0	40	8	8	+	+	-
4	9th month	Stillbirth	Placenta, heart, kidneys, spleen	Echovirus 27	ND	ND	ND	ND	ND	ND	ND
5	9th month	Stillbirth	Placenta, brain, liver, brain, liver,	Coxsackie B ₆	ND	ND	ND	ND	ND	ND	ND

- = negative, ND = not done, CF = Complement fixation, NT = Neutralization test for isolated enterovirus, Toxo = toxoplasmosis, Syph = syphilis, Brucel = brucellosis

Brazil, echovirus and other organs (Fonseca *et al.*, occurrence of fatal infection 11 (Landsman *et al.*, 1980; Modlin, 1986). The study made with a group of patients of maternal infection and its possible damage

pregnant women that had been in Rio de Janeiro, Brazil. The women were collected monthly and two weeks before the end of pregnancy for viral and bacterial cultures, during delivery, for stillbirth cases, tissue samples from placenta and lungs.

and stool samples were inoculated into LLC-MK₂ cells with Melnick's medium and were positive for cytopathogenic effect.

Other were assayed against the complement fixation test to exclude other infections, syphilis, histiocytosis, and glycaemia, and blood pressure

tissue samples. The sections were stained several times with PBS pH 7.2, and isolated viruses for one hour at 4°C. After one hour incubation at 37°C, the sections were stained (Lennette and Schmidt,

four were cut into small pieces and fixed with pH 7.2 cacodylate-sucrose solution, a series of alcohols and embedded in epoxy ultramicrotome. The sections were stained and observed in electron micro-

abortions occurred in the 5th month (one case) and 2.5% by stillbirth. The viruses were isolated from 3 maternal cases such as brain, liver, kidney, and placenta. It has been isolated from 4 cases




Fig. 1.

Specific immunofluorescence in placental tissue confirming infection with Echovirus type 27 (case 4). Scattered foci of immunofluorescence (arrows), magn. 400 \times .




Fig. 2.

Part of foetal hepatocyte infected with Echovirus type 27; dilatation of endoplasmic reticulum in the vicinity of virus-like particles (arrow). The cell cytoplasm is lysed showing aggregates of ribosomes. Bar = 160 nm.

of abortion and stillbirth (Table 1). Echovirus type 27 were isolated from cases 3 and 4.

Enterovirus specific immunofluorescence was observed in three cases of abortion. In two of these children were born with congenital anomalies also observed in all mothers with syphilis, and listeriosis conversions.

The presence of virus was confirmed by immunofluorescence mainly in the villi and chorionic plate in cases 3 and 4 as well as the placenta. It had been investigated whether any alterations were possible in the placenta. We found alterations in the cytotrophoblasts, syncytiotrophoblasts and formation of intervillous spaces (cases 3, and 4).



Part of a cytotrophoblast infected with Echovirus type 27; N = nucleus, Bar = 175 nm.

of abortion and stillbirth and Coxsackie B types 2 and 6 from two cases (Table 1). Echoviruses types 18, 21, and 25, Coxsackie B types 2 and 6 were isolated from cases of placentitis only.


Enterovirus specific serological conversions in mothers were observed in three cases of abortion. In the group where only placentitis occurred and the children were born without abnormalities, serological conversions were also observed in all mother's sera. Rubella, cytomegalovirus, toxoplasmosis, syphilis, and listeriosis serology were performed without positive serological conversions.

The presence of viral antigens of the enteroviruses here involved, was confirmed by immunofluorescence test. It was positive in the placentas mainly in the villi and Langhan's cells (Fig. 1). All placentas from abortion cases as well as the foetal organs, from where enteroviruses were isolated, had been investigated by electron microscopy. The presented morphological alterations were possibly related to virus replication, but no virions were seen. We found assembly of long polyribosomes attached to enlarged elements from the rough endoplasmic reticulum and electrondense aggregates in the cytotrophoblast. Further alterations were enlargement of mitochondria and formation of vacuoles with multiple concentric membranes (Figs. 2, 3, and 4).



1

Infection with Echovirus type 27 (arrows), magn. 400x.



2

Dilatation of endoplasmic reticulum and polyribosomes is lysed showing aggregates of



3

Fig. 3.

Part of a cytotrophoblast cell: arrangement of picorna virus-like crystals in the cytoplasm. N = nucleus, Bar = 175 nm.

Discussion

Despite of the relatively large number of known factors such as chromosomal aberrations, endocrine disturbances, drugs, radiation, physical injuries, malnutrition, and infections, which were well documented, more than 50% of anomalous births remain unexplained (Brown and Arbor, 1968), specially the abortions and stillbirth cases (Mueller *et al.*, 1983). In fact, surveys have estimated that less than 20% of anomalous infants were known to have been exposed to proved teratogenic agents. In Brazil, unfortunately, no statistical approach has been adopted for anomalous births but it can be assumed high (Radis, 1968). The incidence of infectious diseases with contamination of the placenta, specially by haematogenic route is high and includes: bacterial, fungal, viral, and protozootic agents. Concerning virus infections, the following have been detected: rubella, cytomegalovirus, mumps, varicella-zoster, and herpes simplex (Garcia *et al.*, 1985) and as noticed recently, Coxsackie B and echoviruses (Fonseca *et al.*, 1982).

It is well known that the hypothesis on viral aetiology of congenital malformations is based mostly on the studies with rubella and cytomegalovirus. Both are also incriminated as agents of stillbirth and spontaneous abortion




Fig. 4.

Part of a cytotrophoblast cell: picornavirus-like particles in the cytoplasm near to cell protrusions; the cell is lysed and vacuolated
N = nucleus, Va = vacuoles, Bar = 200 nm.

(Hurley, 1981). Efforts have been formed, employing techniques (Hovata *et al.*, 1983) to observe the mothers and foetal tissues present in the children were born after the studied pregnancies with normal babies, it is possible for viruses to affect the placental tissues. Some mothers (from foetal extracts) had antibodies that they experienced the occurrence of the infection would enable the small virus and to reach the foetus are probably the best explanation. The occurrence of an antecedent infection in the mother delivered normal children. On the other hand, the infection is explained by foecal contamination. It has been found in the children that occurred probably by contact from maternal foecal contamination, active virus infection,

One patient had contact with investigation to rubella virus, without seroconversion in the placenta and from the foetus. A spontaneous abortion occurred. The patient presented a respiratory infection discharge followed soon after by a spontaneous abortion and foetal tissues. Molecular biology (Coxsackie virus 33) infection. I was able to detect viral antigens in the placenta. Although the children showed signs of illness during the pregnancy, whether they became ill or not, a nursery outbreak of infection (Gardner *et al.*, 1986).

Due to the importance of the abortion or stillbirth cases, enterovirus infection is considered as a possible virus to infect placental tissues, causing serious

factors such as chromo-
radiation, physical in-
documented, more than
rown and Arbor, 1968),
r *et al.*, 1983). In fact,
ous infants were known
In Brazil, unfortunately,
ealous births but it can
infectious diseases with
rogenic route is high and
agents. Concerning virus
rubella, cytomegalovirus,
pia *et al.*, 1985) and as
seca *et al.*, 1982).
ology of congenital mal-
lla and cytomegalovirus.
ad spontaneous abortion



the cytoplasm near to cell pro-
ted

(Hurley, 1981). Efforts to incriminate additional teratogens were performed, employing the retrospective analysis or prospective investigation (Hovata *et al.*, 1983; Modlin, 1986). This prospective study permitted us to observe the mothers through pregnancy; the data that accumulated so far, provided us some information such as the high incidence of enterovirus particles present in the placenta from pregnancies considered normal. The children were born apparently without any abnormalities. Since 5% of the studied pregnancies were lost and 95% of the mothers delivered apparently normal babies, it is supposed that the placental barrier does not permit viruses to affect the child, although they are able to replicate in the placental tissues. Some mothers from whom viruses were isolated (placental and foetal extracts) had serological conversions to the virus, thus confirming that they experienced virus infection during pregnancy. The widespread occurrence of the infections was subclinical in mothers but the viraemia would enable the small particles (28 nm) to cross the placental barrier easily and to reach the foetus. The age and the degree of placental differentiation are probably the best explanation for the ability of virus to reach the foetus. The occurrence of abortion was noticed in those patients who had experienced infection in the first months of pregnancy while those who had delivered normal children, had enterovirus infection at the end of gestation. On the other hand, the isolation of echovirus from placenta could be explained by foecal contamination during labour. If the same viruses have been found in the children tissues, this would suggest that the infection occurred probably by haemogenic route. The same viruses were also isolated from maternal foecal samples about 20 days before abortion, stressing an active virus infection, followed by viraemia.

One patient had contact with rubella-like exanthem but serological investigation to rubella virus demonstrated her previous immunity against that virus, without seroconversion and Coxsackie B type 2 was isolated from her placenta and from the child autopsy organs (brain and heart). The spontaneous abortion occurred at the third month of pregnancy. Other patient presented a respiratory-like illness with high fever and chills; vagina blood discharge followed soon thereafter with uterine contraction. The mother had a spontaneous abortion and Echovirus 33 was isolated from placenta and foetal tissues. Mother serology confirmed an acute enterovirus (Echovirus 33) infection. Immunofluorescence tests confirmed the presence of viral antigens in the placenta and foetal organs.

Although the children were born apparently normal and did not show signs of illness during their first days of life at the hospital, we are not sure whether they became carriers of the virus or if they were responsible for a nursery outbreak of enteroviruses infections as discussed elsewhere (Modlin, 1986).

Due to the importance of knowledge about the agents that could trigger abortion or stillbirth cases Brown and Karunas, 1977), we call attention to enterovirus infection during pregnancy and to the capacity of such viruses to infect placenta, to replicate in its tissues and eventually to reach the foetus, causing serious lesions to both, specially to the child that would

show, subsequently, several signs of virus infection. Our results also indicate that primaty infection occurring in the beginning of pregnancy eventually could be harmful to the foetus probably resulting in spontaneous abortion. In the fact, as stressed by Modlin (1986), "whereas acute illness in the mother before birth often precedes neonatal Echovirus infection and the infection transmitted vertically from mother to infant may be severe, post-natal transmission of the same serotype results in milder disease".

When completing this manuscript we noticed that also Parvoviruses have been recently isolated from cases of abortion and stillbirth (Brown *et al.*, 1984; Knott *et al.*, 1984; Hall, 1985); thus, the number of viruses related with pregnancy hazard increases without causing apparent clinical illness in the mother, or, as described in some cases, causing a rubella-like illness. The findings for parvovirus association with clinical and congenital diseases are similar to our findings.

Acknowledgements. We are in debt with CNPq and FINEP (Brazil) that awarded us with a grant for this research. We would like to thank to Elizabeth Britto Ferrarezi and Paulo Roberto de Andrade Rios, for their technical help.

References

- Aycock, W. L. (1941): The frequency of Poliomyelitis in pregnancy. *N. Engl. J. Med.* **225**, 405-408.
- Bates, T. (1955): Poliomyelitis in pregnancy, foetus and newborn. *A.M.A. Am. J. Dis. Child.* **90**, 189-195.
- Berkovich, S., and Smithwick, E. M. (1968): Transplacental infection due to Echovirus type 22. *J. Ped.* **72**, 94-96.
- Brown, G. C., and Arbor, A. (1968): Coxsackie virus infections and heart disease. *Am. Health J.* **75**, 145-146.
- Brown, G. C., and Karunas, R. S. (1977): Relationship of congenital anomalies and maternal infection with selected enteroviruses. *Am. J. Epidemiol.* **96**, 207.
- Brown, T., Anand, A., Ritchie, L. D., Clewby, J. P., and Reid, T. S. M. (1984): Intrauterine parvovirus infection associated with hydrops fetalis. *Lancet* **ii**, 1033-1034.
- Fonseca, M. E. F., Garcia, A. G. P., Lagrota, M. H. C., and Wigg, M. D. (1982): The isolation of viruses from placenta and foetal organs. Papers of the 1st Int. Conf. on the *Impact of viral Diseases on the Development of Latin American Countries and the Caribbean Region*, Vol. II, 704-709.
- Garcia, A. G. P. (1974): Maternal herpes-simplex infection causing abortion. Histopathologic study in the placenta. Hospital Rio de Janeiro, 1970, 78, 1267. *Abstracts on Hygiene* **46**, 1092.
- Garcia, A. G. P., Souza Marques, R. L., Lobato, Y. Y., Fonseca, M. E. F., and Wigg, M. D. (1985): Placental pathology in congenital rubella. *Placenta*.
- Greenwald, P. (1958): Malformations caused by necrosis in the embryo illustrated by the effect of selenium compounds on chicken embryos. *Am. J. Pathol.* **34**, 77-103.
- Gregg, N., and McAlister (1941): Congenital cataract following German measles in the mother. *Trans. Ophthalm. Soc. Australia* **3**, 35.
- Hall, S. (1985): Infection with parvovirus during pregnancy. *Br. Med. J.* **290**, 713-714.
- Hanshaw, J. B., and Dudgeon, J. A. (1978): *Viral diseases of the foetus and newborn*; W. B. Saunders Co., London, 3rd ed.
- Holowach, J., Thurston, D. L., and Becker, B. (1957): Congenital defects in infants following and during pregnancy. *J. Ped.* **55**, 689-694.
- Hovata, O., Lipasti, A., Rapola, J., and Karjalainen, O. (1983): Causes of stillbirth: a clinicopathological study of 243 patients. *Brit. J. Obst. Gynecol.* **90**, 691-696.
- Hurley, R. I. (1981): *Virus infections of the fetus and newborn*; W. B. Saunders Co., Philadelphia.
- Waterson (Ed.): *Recent Advances in Paediatrics*, London, 1981.
- Jones, M. J., Kolb, M., and Jones, K. (1981): Echo type 11 infection. *Br. J. Obst. Gynecol.* **88**, 103-104.
- Kleinman, H., Prince, J., and J. T. (1962): Echovirus infection. *J. Clin. Invest.* **41**, 261-269.
- Knott, P. D., Wilpily, G., and J. T. (1984): Parvovirus infection with parvovirus infection. *Br. J. Obst. Gynecol.* **91**, 103-104.
- Landsman, J. B., Grist, N., and J. T. (1981): Parvovirus infection and malformations. *Br. J. Obst. Gynecol.* **88**, 103-104.
- Lenette, E. H., and Schick, B. (1969): Parvovirus infection. *J. Clin. Microbiol.* **20**, 103-104.
- Modlin, J. F. (1980): Perinatal parvovirus infection and chlamydial infections. *Br. J. Obst. Gynecol.* **87**, 103-104.
- Modlin, J. F. (1986): Perinatal parvovirus infection and chlamydial infections. *Br. J. Obst. Gynecol.* **93**, 103-104.
- Mueller, R. F., Sybert, V., and J. T. (1981): Parvovirus infection. *Br. J. Obst. Gynecol.* **88**, 103-104.
- Radis - Reuniao, Análise de 100 casos de infecção por Parvovirus B19. *Revista Brasileira de Ginecologia e Obstetrícia* (1986); **8**, 175-184.
- Rantasalo, I., Penttinen, J., and J. T. (1981): Parvovirus infection after an epidemic period. *Br. J. Obst. Gynecol.* **88**, 103-104.
- Rhodes, A. J. (1980): *Viral infections of the fetus and newborn*; W. B. Saunders Co., London.

Our results also indicate of pregnancy eventually in spontaneous abortion. acute illness in the virus infection and the infant may be severe, post-milder disease".

at also Parvoviruses have stillbirth (Brown *et al.*, number of viruses related apparent clinical illness in ing a rubella-like illness. and congenital diseases

(Brazil) that awarded us with Britto Ferrarezi and Paulo Ro-

nancy. *N. Engl. J. Med.* 225.

orn. *A.M.A. Am. J. Dis. Child.*

fection due to Echovirus type 22.

and heart disease. *Am. Health*

genital anomalies and maternal 207.

4, T. S. M. (1984): Intrauterine 1033-1034.

igg, M. D. (1982): The isolation Int. Conf. on the Impact of viral of the Caribbean Region, Vol. II,

assing abortion. *Histopathologic* 1267. *Abstracts on Hygiene* 46.

oca, M. E. F., and Wigg, M. D.

embryo illustrated by the effect 34, 77-103.

German measles in the mother.

Br. Med. J. 290, 713-714.

of the foetus and newborn; W. B.

ital defects in infants following

3): Causes of stillbirth: a clinico- 691-696.

Hurley, R. I. (1981): Virus infection in pregnancy and the puerperium, pp. 20-55. In A. P. Waterson (Ed.): *Recent Advances in Clinical Virology*, Vol. 3, Churchill Livingstone.

Jones, M. J., Kolb, M., Votava, H. J., Johnson, R. J., and Smith, T. F. (1980): Intrauterine Echo type 11 infection. *Mayo Clin. Proc.* 55 (7-8), 509-512.

Kleinman, H., Prince, J. T., Mathey, W. E., Rosenfield, A. B., Bearmann, J. E., and Syverton, J. T. (1982): Echovirus 9 infection and congenital abnormalities: a negative report. *Pediatrics* 2, 261-269.

Knott, P. D., Wilply, G. A. C., and Anderson, M. J. (1984): Serologically proved intrauterine infection with parvovirus. *Br. Med. J.* 289, 1660.

Landsman, J. B., Grist, N. R., and Ross, C. A. C. (1964): Echo 9 virus infection and congenital malformations. *Br. J. Prev. Soc. Med.* 18, 152-156.

Lennette, E. H., and Schmidt, N. J. (1979): Diagnostic procedures for viral, rickettsial and chlamydial infections. *Am. Public Health Association*, Washington, D.C. 20005, 5th edition.

Modlin, J. F. (1986): Perinatal echovirus infection: insights from a literature review of 61 cases of serious infection and 16 outbreak nurseries. *Rev. infect. Dis* 8 (6), 818-926.

Mueller, R. F., Sybert, V. P., Johnson, J., Brown, Z. A., and Wei Jen Chen (1983): Evaluation of a protocol for post-mortem examination of stillbirths. *N. Engl. J. Med.* 309 (10), 586-590.

Radis - Reuniao, Análise e Difusao de Informacao sobre Saúde. Escola Nacional de Saúde Pública. *Fiocruz* (1986): 9.

Rantasalo, I., Penttineu, K., Saxen, L., and Ojala, A. (1960): Echo 9 virus antibody status after an epidemic period and possible teratogenic effect of the infection. *Ann. Paediatr. Fenn.* 6, 175-184.

Rhodes, A. J. (1980): Virus infection and congenital malformations, In: *1st Conf. Congenital Malformations*, London.