




Correction to: Expression of *O*-glycosylated oncofetal fibronectin in alternatively activated human macrophages

Marcos Andre Rodrigues da Costa Santos¹ · Jhenifer Santos dos Reis¹ · Carlos Antonio do Nascimento Santos¹ · Kelli Monteiro da Costa¹ · Pedro Marçal Barcelos¹ · Karen Queiroz de Oliveira Francisco¹ · Pedro Antônio Guimarães Notaroberto Barbosa¹ · Emanuelle Damasceno Souza da Silva¹ · Celio Geraldo Freire-de-Lima¹ · Alexandre Morrot^{2,3} · Debora Decote-Ricardo⁴ · Israel Diniz-Lima¹ · Jose Osvaldo Previato¹ · Lucia Mendonca-Previato¹ · Leonardo Marques da Fonseca¹ · Leonardo Freire-de-Lima¹ 

Published online: 10 November 2022

© Springer Science+Business Media, LLC, part of Springer Nature 2022

Correction to: Immunologic Research

<https://doi.org/10.1007/s12026-022-09321-9>

In the originally published version of this article, the authors have noticed mistakes during figure production, where Supplementary Fig. 3 was incomplete, missing Western blot data. The complete Supplementary Fig. 3A,B,C is displayed in the next page.

The original article has been corrected.

Marcos Andre Rodrigues da Costa Santos, Jhenifer Santos dos Reis, and Carlos Antonio do Nascimento Santos contributed equally to this work.

The original article can be found online at <https://doi.org/10.1007/s12026-022-09321-9>.

✉ Leonardo Marques da Fonseca
lfonseca@biof.ufrj.br

✉ Leonardo Freire-de-Lima
leolima@biof.ufrj.br


¹ Instituto de Biofísica Carlos Chagas Filho, Laboratório de Biologia Celular de Glicoconjugados, Universidade Federal Do Rio de Janeiro, Rio de Janeiro, RJ 21941-902, Brazil

² Fiocruz, Laboratório de Imunoparasitologia, Instituto Oswaldo Cruz, Rio de Janeiro, RJ 21040-360, Brazil

³ Faculdade de Medicina, Universidade Federal Do Rio de Janeiro, Rio de Janeiro, RJ 21941-902, Brazil

⁴ Departamento de Microbiologia E Imunologia Veterinária, Instituto de Veterinária, Universidade Federal Rural Do Rio de Janeiro, Rio de Janeiro, RJ 23890-000, Brazil

Supplementary Fig. 3 Analysis of GalNAc-T6 expression in THP-1 derived macrophages by qPCR and Western blot. After differentiation from Mon ϕ to M ϕ with PMA, cells were classically or alternatively activated as detailed in the material and methods section. mRNA and protein levels for pp-GalNAc-T6 in both M1-M ϕ and M2-M ϕ were monitored by real time PCR (A) and Western blot (B and C), respectively. Graphs represent three individual experiments. n.s., not significant in relation to M ϕ .



Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.