

Journal Pre-proof

SARS-CoV-2 epidemic in Brazil: how variants displacement have driven distinct epidemic waves

Luiz Carlos Junior Alcantara , Elisson Nogueira , Gabriel Shuab ,
Stephane Tosta , Hegger Fristch , Victor Pimentel ,
Jayme A. Souza-Neto , Luiz Lehmann Coutinho ,
Heidge Fukumasu , Sandra Coccuzzo Sampaio ,
Maria Carolina Elias , Simone Kashima , Svetoslav Nanev Slavov ,
Massimo Ciccozzi , Eleonora Cella , José Lourenco ,
Vagner Fonseca , Marta Giovanetti



PII: S0168-1702(22)00112-5
DOI: <https://doi.org/10.1016/j.virusres.2022.198785>
Reference: VIRUS 198785

To appear in: *Virus Research*

Received date: 8 March 2022
Revised date: 11 April 2022
Accepted date: 12 April 2022

Please cite this article as: Luiz Carlos Junior Alcantara , Elisson Nogueira ,
Gabriel Shuab , Stephane Tosta , Hegger Fristch , Victor Pimentel , Jayme A. Souza-Neto ,
Luiz Lehmann Coutinho , Heidge Fukumasu , Sandra Coccuzzo Sampaio , Maria Carolina Elias ,
Simone Kashima , Svetoslav Nanev Slavov , Massimo Ciccozzi , Eleonora Cella ,
José Lourenco , Vagner Fonseca , Marta Giovanetti , SARS-CoV-2 epidemic in Brazil: how
variants displacement have driven distinct epidemic waves, *Virus Research* (2022), doi:
<https://doi.org/10.1016/j.virusres.2022.198785>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 Published by Elsevier B.V.

SARS-CoV-2 epidemic in Brazil: how variants displacement have driven distinct epidemic waves

Luiz Carlos Junior Alcantara^{1,2,*^} luiz.alcantara@ioc.fiocruz.br, Elisson Nogueira^{2,*}, Gabriel Shuab^{1,*}, Stephane Tosta^{2,*}, Hegger Fristch², Victor Pimentel³, Jayme A. Souza-Neto⁴, Luiz Lehmann Coutinho⁵, Heidge Fukumasu⁶, Sandra Coccuzzo Sampaio⁷, Maria Carolina Elias⁷, Simone Kashima⁸, Svetoslav Nanev Slavov⁸, Massimo Ciccozzi⁹, Eleonora Cella¹⁰, José Lourenco^{11,12}, Vagner Fonseca^{13,14,^} vagner.fonseca@gmail.com, Marta Giovanetti^{1,2,15,^} giovanetti.marta@gmail.com

¹Laboratório de Flavivírus, Instituto Oswaldo Cruz, Fundação Oswaldo Cruz, Rio de Janeiro, Brazil

²Laboratório de Genética Celular e Molecular, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais, Brazil

³Global Health and Tropical Medicine (GHTM), Instituto de Higiene e Medicina Tropical/Universidade Nova de Lisboa (IHMT/UNL), Lisbon, Portugal

⁴São Paulo State University (UNESP), School of Agricultural Sciences, Botucatu, Brazil

⁵University of São Paulo, Centro de Genômica Funcional da ESALQ, Piracicaba, SP, Brazil

⁶Department of Veterinary Medicine, School of Animal Science and Food Engineering, University of Sao Paulo, Pirassununga, Sao Paulo, Brazil

⁷Butantan Institute, São Paulo, Brazil

⁸University of São Paulo, Ribeirão Preto Medical School, Blood Center of Ribeirão Preto, Ribeirão Preto, SP, Brazil

⁹Medical Statistic and Molecular Epidemiology Unit, University of Biomedical Campus, Rome, Italy

¹⁰Burnett School of Biomedical Sciences, University of Central Florida, Orlando, FL, USA

¹¹Department of Zoology, Peter Medawar Building, University of Oxford, Oxford, UK

¹²Biosystems and Integrative Sciences Institute (BioISI), Universidade de Lisboa, Portugal

¹³Organização Pan-Americana da Saúde/Organização Mundial da Saúde, Brasília, Distrito Federal, Brazil

¹⁴KwaZulu-Natal Research Innovation and Sequencing Platform (KRISP), School of Laboratory Medicine and Medical Sciences, College of Health Sciences, University of KwaZulu-Natal, Durban, South Africa;

¹⁵Department of Science and Technology for Humans and the Environment, University of Campus Bio-Medico di Roma, Rome, Italy.

^Corresponding Author.

*Denote equal contribution

Highlights

- Brazil ranks as third in terms of total number of reported SARS-CoV-2 cases globally.
- Here we show how variants displacement have driven distinct epidemic waves in Brazil.
- Genomic effort is pivotal to follow the evolution of this unpredictable virus.

Abstract

Brazil ranks as third in terms of total number of reported SARS-CoV-2 cases globally. The COVID-19 epidemic in Brazil was characterised by the co-circulation of multiple variants as a consequence of multiple independent introduction events occurring through time. Here, we describe the SARS-CoV-2 variants that are currently circulating and co-circulating in the country, with the aim to highlight which variants have driven the different epidemic waves. For this purpose, we retrieved metadata information of Coronavirus sequences collected in Brazil and available at the GISAID database. SARS-CoV-2 lineages have been identified along with eleven variants, labelled as VOCs (Alpha, Gamma, Beta, Delta and Omicron) VOIs (Lambda and Mu) VUMs (B.1.1.318) and FMVs (Zeta, Eta and B.1.1.519). Here we show that, in the Brazilian context, after 24 months of sustained transmission and evolution of SARS-CoV-2, local variants (among them the B.1.1.28 and B.1.1.33) were displaced by recently introduced VOCs firstly with the Gamma, followed by Delta and more recently Omicron. The rapid spread of some of those VOCs (such as Gamma and Omicron) was also mirrored by a large increase in the number of cases and deaths in the country. This in turn reinforces that, due to the emergence of variants that appear to induce a substantial evasion against neutralizing antibody response, it is important to strengthen genomic effort within the country and how vaccination still remains a critical process to protect the vulnerable population, still at risk of infection and death.

Keywords

SARS-CoV-2, variants, Brazil, surveillance

Introduction

The Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), an emerging RNA Betacoronavirus first detected in China by the end of 2019, is considered a major public health threat. Since the onset of the COVID-19 pandemic, SARS-CoV-2 variants have been emerging and circulating around the world and have made major contributions to the recurring epidemic waves that occurred asynchronously in different regions through time. During late 2020, the emergence of variants that posed an increased risk to the global public health prompted the implementation of major monitoring programs in order to inform the ongoing response to the COVID-19 pandemic. This in turn yielded the generation as of the end of February 2022 of more than 8 million genomes through sequencing-based surveillance¹.

Currently, lineages that may require public health actions have been classified by the WHO as variant of concern (VOC), variant of interest (VOI), variant under monitoring (VUM) and as formerly monitored variant (FMV)².

The classification of variants occurs dynamically according to the evolution of the virus and follows the Greek letter system that denominates variants in a non-stigmatising manner, as advised by WHO².

VUMs generally require epidemiological or phenotypic assessment to investigate suspicious genetic changes that may enhance viral fitness. If it is predicted or established that defined genetic changes may affect diagnostic and therapeutic strategies, immune escape, transmissibility, and disease severity, the investigated VUM starts to be considered as a VOI. Additionally, when genetic properties are associated with increased transmissibility and virulence, detrimental change in clinical presentation and a decrease in effectiveness of public health measures, the VOI starts to be classified as a VOC².

As the spectrum of knowledge regarding the variants may rapidly evolve, variants can further be reassigned as FMVs. This happens when those variants start to be associated with reduced public health significance due to low frequencies of circulation, undetected impact in a long-term period and when they are no longer associated with factors of concern².

In Brazil, the COVID-19 epidemic had huge impact. According to the National health report Brazil reported more than 27 million confirmed cases and a death toll exceeding 640 thousand by the end of February of 2022, making it one of the countries hardest hit by COVID-19 pandemic^{3,4}.

In this study we provide insights regarding the spread of SARS-CoV-2 variants in the Brazilian territory, highlighting how the replacement of several variants of concern have dictated the different epidemic waves in the country.

Materials and methods

We retrieved metadata information of Brazilian SARS-CoV-2 sequences available at the GISAID¹ database collected up to February 19th, 2022. To ensure the quality of the data analysed in this study

and to guarantee the highest possible accuracy of the obtained results, only genomes >29,000bp and <1% of ambiguities with a variant assignment provided by the Phylogenetic Assignment of Named Global Outbreak Lineages (PANGOLIN)⁵, were considered (n=111,626). For convenience, the geographical locations were aggregated in five Brazilian macro regions: North, Northeast, Southeast, Midwest and South. North macro region includes the states of: Acre, Amapá, Amazonas, Pará, Rondônia, Roraima, Tocantins. Northeast macro region includes the states of: Alagoas, Bahia, Ceará, Maranhão, Paraíba, Pernambuco, Piauí, Rio Grande do Norte, Sergipe. Southeast macro region includes the states of: Rio de Janeiro, Espírito Santo and Minas Gerais. Midwest macro region include the states of: Goiás, Federal District, Mato Grosso and Mato Grosso do Sul. South macro region includes the states of: Parana, Santa Catarina and Rio Grande do Sul.

In addition, daily cases of SARS-CoV-2 in Brazil were retrieved from the Official National repository available at covid.saude.gov.br. The National network releases daily updates on the number of confirmed new cases, deaths, and recoveries, with a breakdown by states and regions.

Results

As a consequence of efforts to generate genomic data that would contribute to the mitigation of the pandemic, by the end of February 2022, a total number of 240 different lineages had been identified in Brazil. Eleven of them were assigned as: VOCs (Alpha, Gamma, Beta, Delta and Omicron); VOIs (Lambda and Mu); VUMs (B.1.1.318) and FMVs (Zeta, Eta and B.1.1.519). Ancestral relationships among these variants and main lineages identified in Brazil are represented as a diagram in (Fig. 1).

The Brazilian epidemic has been characterized by three distinct epidemic waves, causing more than 27 million cases and 670 thousand deaths as of 19th February 2022 (Fig. 2 and Fig. S1)⁶. The first wave from February 2020 to November 2020 was characterized by the co-circulation of different lineages because of multiple introduction events occurring through time (Fig. 2A)⁴. The second epidemic wave (December 2020 up to December 2021) was fuelled by the emergence and circulation of several VUMs, such as P.2 (i.e., Zeta), and VOCs such as Gamma (i.e., P.1), which started to be detected from January 2021 (Fig. 2A and Fig. S1). We detected several additional VOCs, VOIs, VUMs and FMVs sporadically during this wave period, including the Alpha, Mu and Eta, C.1.2, B.1.1.318, Lambda and B.1.1.519 variants from January 2021 onwards, but they remained at a low frequency nationally and were not associated with a significant resurgence of cases in any regions (Fig. S2). In April 2021, the Delta VOC started to be detected in the country. This VOC was able to displace the Gamma variant becoming the dominant viral circulating variant in the National scenario by the end of October 2021 (Fig. 2A and Fig. S1)⁷.

The months following the detection of this emerging VOC in Brazil, between September 2021 and December 2021, were marked by lower levels of transmission as indicated by a low incidence of reported COVID-19 cases and deaths (**Fig. 2**). It is nonetheless challenging to identify the exact cause of this trend, but it is likely that immunity acquired by prior infection with previous variants (e.g. Gamma) together with a successful vaccination coverage (more than 50%) played a role in the declining numbers of reported cases and deaths in the country (**Fig. 2**)⁷.

After the emergence of Omicron in South Africa and Botswana, in late November 2021, the first Omicron imported cases was also detected in Brazil^{8,9}. Since then, we observed a striking growth in the prevalence rate of this emerging variant from early January 2022. As already described before, it can be noted a clear trend of Delta's replacement by Omicron (**Fig. 2A**) at the national and international level^{9,10}. The Omicron variant (B.1.1.529) raised a world alert due to its constellation of mutations (30 mutations^{9,11}), most of them located in the spike protein^{11,12} and likely-associated with increased risk of infection¹³. Several Omicron's sublineages (BA.1, BA.2, BA.3, among others) have been already described¹⁴, with BA.3 as the least prevalent. Studies have shown that BA.2 has a growth advantage over BA.1, which currently remains the most common Omicron sublineage reported worldwide¹⁴. This in turn reinforce how crucial appear to be following their realtime evolution worldwide.

A month after the first identified Omicron case in the country, this emerging variant was associated with sustained transmission within all Brazilian regions and was also associated with an initial increase in COVID-19 cases and deaths (**Fig 2 and Fig. S1**). It is likely that the Omicron mutations allowed to better evade immune protection while spreading faster than any prior known variant⁸. Despite this, it was observed that the Omicron's wave had a sharp peak and a swift decline, likely reflecting the successful impact of the vaccination programme together with the capacity of this unpredictable and very transmissible virus to quickly affect the proportion of susceptible population.

Discussion

Brazil has historically faced one of the worst COVID-19 scenarios worldwide and made South America the epicentre of the SARS-CoV-2 epidemic. Different sequencing efforts have been employed by Brazilian states thus resulting in differing numbers of available genomes for each state/region (totalizing n=113,681 complete genome sequences available as of 19th February 2022), thus struggling our ability to describe in detail the progression of the epidemic in the country. Considering that since the beginning of the epidemic sustained lineage replacements over time have been observed^{4,7}, our study reinforces how important it is to strengthen genomic monitoring in order to follow local time evolution of this viral pathogen.

In this study by combining epidemiological and genomic data we show that after 24 months of persistent transmission and evolution of SARS-CoV-2, local variants (among them the B.1.1.28 and the B.1.1.33) were subsequently displaced by recently described VOCs, firstly by Gamma, followed by Delta and Omicron, and how the rapid spread of some of those variants also dictated a large increase in the number of cases and deaths in the country. Understanding why cases are rising and falling is crucial for implementing adequate control measures.

Omicron drove a third wave of SARS-CoV-2 cases and deaths in Brazil, as it did rapidly worldwide. Additionally, the recent identification of Omicron's sub lineages (BA.1, BA.2, BA.3, among others) raises concern regarding their potential impact on the national and international health systems. Close monitoring of the Omicron's sublineages evolution appears to be necessary to better understand its transmissibility and the capacity of this variant to evade post-infection and vaccine-elicited cases. Our results finally highlight the need of more public investments to strengthen the genomic capacity across the country.

Author contribution

Conceptualization: LCJA, and MG; Curated metadata: EN, GS; ST.; Analysed the data: LCJA, EN, GS; ST., VF., and MG.; Helped with study design and data interpretation: LCJA; EN, GS, ST, HF, VP, SA, JASN, LLC, HF, SCS, MCE, SK, SNS, EC, MC, JL, VF, and MG ; Wrote the initial manuscript, which was reviewed by all authors: EN, ST, HF and MG.

Supplementary information

Supplementary Figure 1. Dynamics of the SARS-CoV-2 epidemic in each Brazilian microregion showing the number of daily COVID-19 deaths and the progression in the proportion of circulating variants in the country over time, showing the rapid replacement of several VOCs (Gamma, Delta and Omicron).

Supplementary Figure 2. Frequency and distribution of SARS-CoV-2 VOCs, VOIs, VUMs and FMVs in Brazil.

Declaration of Competing Interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Funding statement

This work was supported in part through National Institutes of Health USA grant U01 AI151698 for the United World Arbovirus Research Network (UWARN), the CRP- ICGEB RESEARCH GRANT 2020 Project CRP/BRA20-03, Contract CRP/20/03, the Oswaldo Cruz Foundation VPGDI-027-FIO-20-2-2-30, and the Brazilian Ministry of Health (SCON2021-00180). MG is supported by Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro -FAPERJ.

Journal Pre-proof

References

1. Khare, S., et al (2021) GISAID's Role in Pandemic Response. *China CDC Weekly*, 3(49): 1049-1051. doi: 10.46234/ccdcw2021.25 PMID: 8668406
2. World Health Organization. Tracking sars-COV-2 variants. World Health Organization. Retrieved February 21, 2022, from <https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/>
3. World Health Organization. Retrieved February 21, 2022, from <https://covid19.who.int/>
4. Giovanetti M., et al., Genomic epidemiology reveals how restriction measures shaped the SARSCoV- 2 epidemic in Brazil. *MedRxiv*, Oct 2021. doi: <https://doi.org/10.1101/2021.10.07.21264644>
5. O'Toole, Á., Scher, E., Underwood, A., Jackson, B., Hill, V., McCrone, J. T., Colquhoun, R., Ruis, C., Abu-Dahab, K., Taylor, B., Yeats, C., Du Plessis, L., Maloney, D., Medd, N., Attwood, S. W., Aanensen, D. M., Holmes, E. C., Pybus, O. G., & Rambaut, A. (2021). Assignment of epidemiological lineages in an emerging pandemic using the pangolin tool. *Virus Evolution*, [veab064]. <https://doi.org/10.1093/ve/veab064>
6. COVID-19 Situation Reports - PAHO/WHO | Pan American Health Organization. <https://www.paho.org/en/covid-19-situation-reports>.
7. Giovanetti M. et al., Replacement of the Gamma by the Delta variant in Brazil: impact of lineage displacement on the ongoing pandemic. *MedRxiv*, Dec, 2021. doi: <https://doi.org/10.1101/2021.12.27.21268309>.
8. Viana R. et al. Rapid epidemic expansion of the SARS-CoV-2 Omicron variant in southern Africa. *Nature*, doi:doi: 10.1038/d41586-021-03832-5 (2021).
9. Campos Soares G et al. (2022). SARS-CoV-2 epidemic in Brazil: how variants displacement have driven distinct epidemic waves Genomic monitoring unveils a high prevalence of SARS-CoV-2 Omicron variant in vaccine breakthrough cases. *MedRxiv*. doi: <https://doi.org/10.1101/2022.02.16.22271059>
10. Callaway, E. (2021). Heavily mutated Omicron variant puts scientists on alert. In *Nature* (Vol. 600, Issue 7887, pp. 21–21), <https://doi.org/10.1038/d41586-021-03552-w>

11. Saito, A., Nasser, H., Uriu, K., Kosugi, Y., Irie, T., Shirakawa, K., Sadamasu, K., Kimura, I., Ito, J., Wu, J., Ozono, S., Tokunaga, K., Butlertanaka, E. P., Tanaka, Y. L., Shimizu, R., Shimizu, K., Fukuhara, T., Kawabata, R., Sakaguchi, T., ... Sato, K. (2021). SARS-COV-2 spike P681R mutation enhances and accelerates viral fusion. <https://doi.org/10.1101/2021.06.17.448820>
12. Gong, S. Y., Chatterjee, D., Richard, J., Prévost, J., Tausin, A., Gasser, R., Bo, Y., Vézina, D., Goyette, G., Gendron-Lepage, G., Medjahed, H., Roger, M., Côté, M., & Finzi, A. (2021). Contribution of single mutations to selected SARS-COV-2 emerging variants Spike Antigenicity. <https://doi.org/10.1101/2021.08.04.455140>
13. Wolter, N., Jassat, W., Walaza, S., Welch, R., Moultrie, H., Groome, M., Amoako, D. G., Everatt, J., Bhiman, J. N., Scheepers, C., Tebeila, N., Chiwandire, N., du Plessis, M., Govender, N., Ismail, A., Glass, A., Mlisana, K., Stevens, W., Treurnicht, F. K., ... Cohen, C. (2021). Early assessment of the clinical severity of the SARS-COV-2 omicron variant in South Africa. <https://doi.org/10.1101/2021.12.21.21268116>
14. World Health Organization. (2022) Statement on Omicron sublineage BA.2. Available at: <https://www.who.int/news/item/22-02-2022-statement-on-omicron-sublineage-ba.2>

Figure legends




Figure 1. Diagram of variants detected in Brazil. The diagram represents the evolutionary relation of the twelve VOCs (alpha, beta, gamma, delta, omicron), VOIs (mu, lamda), VUMs (B.1.1.318) and FMVs (zeta, eta and B.1.1.519 detected in Brazil since the beginning of the epidemic until the February 19th, 2022. Colours indicate WHO variant classification. Main ancestor lineage is B.1. Lineage B.1.1.28 had a crucial impact leading to variants that presented high frequency lately in Brazilian genomes (Gamma and Zeta).




Figure 2. Dynamics of the SARS-CoV-2 epidemic in Brazil. A) Number of daily COVID-19 cases and the vaccination rates in time; B) Progression in the proportion of circulating variants in Brazil over the first second and third waves of infection, showing the rapid replacement of different VOCs throughout time additionally showing the number of daily COVID-19 deaths.