

NADPH Phagocyte Oxidase Knockout Mice Control *Trypanosoma cruzi* Proliferation, but Develop Circulatory Collapse and Succumb to Infection

Helton C. Santiago^{1, *a}, Claudia Z. Gonzalez Lombana^{1, *b}, Juan P. Macedo¹, Lara Utsch¹, Wagner L. Tafuri², Maria José Campagnole-Santos³, Rosana O. Alves⁴, José C. F. Alves-Filho⁵, Alvaro J. Romanha⁴, Fernando Queiroz Cunha⁵, Mauro M. Teixeira¹, Rafael Radi^{6,8§}, Leda Q. Vieira^{1,7*¶}

1 Departamento de Bioquímica e Imunologia, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais, Brazil, **2** Departamento de Patologia Geral, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais, Brazil, **3** Departamento de Fisiologia e Biofísica, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais, Brazil, **4** Centro de Pesquisas René Rachou, Fiocruz, Belo Horizonte, Minas Gerais, Brazil, **5** Departamento de Bioquímica e Imunologia, Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, Brazil, **6** Departamento de Bioquímica, Universidad de la República, Montevideo, Uruguay, **7** Núcleo de Pesquisas em Ciências Biológicas, Universidade Federal de Ouro Preto, Ouro Preto, Minas Gerais, Brazil, **8** Center for Free Radical and Biomedical Research, Universidad de la República, Montevideo, Uruguay

Abstract

NO is considered to be a key macrophage-derived cytotoxic effector during *Trypanosoma cruzi* infection. On the other hand, the microbicidal properties of reactive oxygen species (ROS) are well recognized, but little importance has been attributed to them during *in vivo* infection with *T. cruzi*. In order to investigate the role of ROS in *T. cruzi* infection, mice deficient in NADPH phagocyte oxidase (gp91^{phox} / or *phox* KO) were infected with Y strain of *T. cruzi* and the course of infection was followed. *phox* KO mice had similar parasitemia, similar tissue parasitism and similar levels of IFN- γ and TNF in serum and spleen cell culture supernatants, when compared to wild-type controls. However, all *phox* KO mice succumbed to infection between day 15 and 21 after inoculation with the parasite, while 60% of wild-type mice were alive 50 days after infection. Further investigation demonstrated increased serum levels of nitrite and nitrate (NOx) at day 15 of infection in *phox* KO animals, associated with a drop in blood pressure. Treatment with a NOS2 inhibitor corrected the blood pressure, implicating NOS2 in this phenomenon. We postulate that superoxide reacts with NO *in vivo*, preventing blood pressure drops in wild type mice. Hence, whilst superoxide from phagocytes did not play a critical role in parasite control in the *phox* KO animals, its production would have an important protective effect against blood pressure decline during infection with *T. cruzi*.

Citation: Santiago HC, Gonzalez Lombana CZ, Macedo JP, Utsch L, Tafuri WL, et al. (2012) NADPH Phagocyte Oxidase Knockout Mice Control *Trypanosoma cruzi* Proliferation, but Develop Circulatory Collapse and Succumb to Infection. PLoS Negl Trop Dis 6(2): e1492. doi:10.1371/journal.pntd.0001492

Editor: Diane McMahon-Pratt, Yale School of Public Health, United States of America

Received: December 13, 2010; **Accepted:** December 9, 2011; **Published:** February 14, 2012

Copyright: © 2012 Santiago et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: FUNDEP was crucial for logistic support. This work was supported by grants from Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq473597/2006-3 and 304776/2009-2), CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior) and INCT de Processos Redox em Biomedicina-Redoxoma (CNPq/FAPESP/CAPE5573530/2008-4). HCS, CZGL, JPM, LU, WLT, MJCS, AJR, FQC, MMT and LQV are CNPq fellows. RR was supported by grants from the National Institutes of Health (1R01AI095173-01), Howard Hughes Medical Institute and Comisión Sectorial de Investigación Científica (CSIC)/Universidad de la República, Uruguay. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: lqviera@icb.ufmg.br

¶ These authors contributed equally to this work.

§ These authors also contributed equally to this work.

^a Current address: Laboratory of Parasitic Diseases, National Institutes of Health, Bethesda, Maryland, United States of America

^b Current address: Department of Pathobiology, School of Veterinary Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, United States of America

Introduction

For a long time, reactive oxygen species (ROS) were considered the main anti-microbial radical produced by the immune system, playing a role against bacterial, fungal and protozoa infections. After the discovery of nitric oxide (NO), NO found to play a major role in host defense, especially against protozoan parasites. A role against *Toxoplasma* [1,2], *Plasmodium* [3] and *Leishmania* [4,5] infections was still attributed to ROS, albeit in some cases this role remains a matter of debate [6,7,8,9].

Since NO was found to be one of the most important IFN- γ -induced anti-parasitic mechanisms, the studies about its role in

different diseases was intensified. The advent of gene knockout (KO) technology allowed the dissection of the real extent of NO involvement in parasitic diseases. NO was found to be crucially important in a variety of infections [10,11], however, NOS2-deficient animals are less susceptible than *ifn- γ* KO to most microorganisms studied [12,13,14,15,16]. So, the search for other mechanisms of host resistance induced by IFN- γ started, and the interest in ROS warmed up again.

Trypanosoma cruzi is an intracellular parasite associated with high morbidity during both acute and chronic phases of infection. Resistance to this parasite is mostly driven by IFN- γ . This cytokine mediates the control of parasite proliferation in tissues

Author Summary

When pathogens enter their hosts, they are fought by several resistance strategies, including capture by phagocytes and the production of pathogen-toxic molecules. Nitric oxide, a free radical, has been extensively studied as one of these toxic molecules that successfully mediates intracellular parasite killing, including *Trypanosoma cruzi*, the protozoan parasite that causes Chagas' disease. On the other hand, reactive oxygen species also mediate resistance to several pathogens, mainly bacterial. In this study, we addressed the role of reactive oxygen species in the resistance to *T. cruzi* using gene-deficient mice, a species which phagocytes lack the ability to produce (*phox*^{-/-} mice). We found that phagocyte-derived reactive oxygen species are not critical to mediate resistance to parasite in the knock-out animals. However, *phox*^{-/-} mice presented higher mortality and lower blood pressure due to infection with *T. cruzi* than non-deficient mice. The blood pressure was restored to normal by an inhibitor of nitric oxide synthesis by phagocytes. We hypothesize that superoxide (one of the oxygen reactive species) controls blood pressure during infection with *T. cruzi*, by reacting with nitric oxide and preventing its action on blood vessels.

and blood in a NOS2-dependent way. However, NO may not be necessary for host resistance to *T. cruzi* infection when less virulent strains are used [13]. In addition, previously published data suggest that NOS2 deficient mice exhibit delayed mortality when compared to *ifn-γ* KO mice [13,14], denoting an additional effector mechanism involved in *T. cruzi* immune resistance. Further studies suggested IFN-γ-induced p47GTPase LRG-47 as one major factor of resistance to *T. cruzi* infection along with NO [17,18]. Although there is convincing evidence for the effects of ROS-induced damage to *T. cruzi* *in vitro* [19,20], the role of these reactive species *in vivo* has not yet been addressed.

In vitro, *T. cruzi* is readily phagocytosed by macrophages and triggers respiratory burst [19,21]. However, production of ROS alone is not sufficient to kill parasites inside these cells [20,21], and activation by IFN-γ, induction of NOS2 and production of NO are required [20,21,22]. In the infected macrophage, NO reacts with superoxide yielding peroxynitrite [21], which is a powerful oxidant and seems to be the main effector molecule against *T. cruzi* [19]. Peroxynitrite is more efficient to kill *T. cruzi* epimastigotes *in vitro* than superoxide or NO alone [19]. Moreover, evidence of peroxynitrite production during *in vitro* and *in vivo* infection with *T. cruzi* is available, as nitrated proteins are found both in macrophages and in mouse and human tissues [23,24]. Indeed, it has just been reported that internalized trypomastigotes in activated macrophages are killed by peroxynitrite-dependent mechanisms [21]. The importance of nitro-oxidative mechanisms is underscored by the finding that virulent *T. cruzi* strains, which naturally have high peroxiredoxin levels [25], and strains overexpressing peroxiredoxins [21,26] are protected from peroxynitrite and macrophage-dependent nitro-oxidative killing (peroxiredoxins readily decompose peroxynitrite). Albeit nitration of proteins *in vivo* may be achieved independently of peroxynitrite, it is still dependent on the production of superoxide and NO [23,24,27]. Hence, parasite damage is dependent not only on ·NO, but on both superoxide and nitric oxide.

In order to investigate the contribution of ROS in resistance to *T. cruzi* infection, mice deficient in the gp91^{phox} (*phox* KO) subunit of NADPH oxidase, a model for chronic granulomatous disease

[28], were used. These animals fail to produce ROS in endothelial cells, causing a defect in endothelium-derived relaxation of arteries [29,30], and in phagocytic cells, leading to deficient resolution of bacterial and fungal infections [28]. Although these animals were found somewhat more susceptible to *Leishmania donovani* [5], their susceptibility to *L. major* is still a matter of debate [4,6]. In the present study, *phox* KO mice were found to succumb to infection with *T. cruzi*, despite adequate control of parasite replication. The immunological and physiological functions of ROS in such model were investigated.

Methods

Ethics statement

The procedures used in this study were approved by the Animal Ethics committee at the Universidade Federal de Minas Gerais, protocol number 031/09. All care was taken to minimize animal suffering.

Animals

Inbred C57BL/6 (WT) mice (males and females, 4–6 week old) were used as controls (CEBIO, Instituto de Ciências Biológicas, UFMG, Belo Horizonte, MG, Brazil). Animals were kept in a conventional animal facility at controlled temperature, light/dark cycles and environmental barriers. The gp91^{phox}-deficient (*phox* KO) [28] and IFN-γ-deficient (*ifn-γ* KO) [31] mice, both in C57BL/6 background, were purchased from The Jackson Laboratories (Bar Harbor, ME, USA) and bred under specific pathogen free conditions at the Gnotobiology Laboratory, Departamento de Bioquímica e Imunologia, ICB, UFMG.

Parasite, infection, cytokines and serum NOx measurements

T. cruzi (Y strain) was maintained by weekly passage in Swiss mice. For *in vivo* experimental infections, mice were inoculated i.p. with 1000 blood-stage trypomastigotes. The parasitemia was evaluated by counting parasites in 5 μL of blood drawn from the tail vein [32]. Mortality of infected mice was monitored daily. Spleen cell cultures were performed as previously described [32]. Briefly, splenocytes from infected mice were obtained on day 10 after infection, and cultured at 5 × 10⁶ cells/ml, in 24-well plates, with RPMI 1640 supplemented with 10% FCS, 2 mM L-glutamine, 0.05 mM 2-mercapto-ethanol, 100 U/ml penicillin, and 100 μg/ml streptomycin. Cultures were maintained at 37°C in 5% CO₂ atmosphere. Supernatants were harvested 72 hours later for TNF and IFN-γ measurements. Mice were bled on days 0, 10 and 15 after infection and the level of serum cytokines was evaluated. IFN-γ and TNF were measured as described previously using specific ELISA kits (R&D Systems, Minneapolis, MN, USA) following the manufacturer's protocol. Nitrate was reduced to nitrite in lipid-free serum with nitrate reductase and measured by the Griess colorimetric reaction [33]. ELISA and immunohistochemistry for 3-nitrotyrosine (or nitrated proteins) was performed as previously described [24].

Quantification of parasite tissue loads and *nos2* mRNA expression by real-time PCR or real-time RT-PCR

Real-time PCR for parasite quantification was performed as described previously [34] with minor modifications. Briefly, on different days after infection, heart, spleen, and liver were digested with proteinase K, followed by a phenol-chloroform-isoamyl alcohol affinity extraction. Real-time PCR using 50 ng of total DNA was performed on an ABI PRISM 7900 sequence detection

system (Applied Biosystems) using SYBR Green PCR Master Mix, according to the manufacturer's recommendations. The equivalence of host DNA in the samples was confirmed by measurement of genomic IL-12p40 PCR product levels in the same samples. Purified *T. cruzi* DNA (American Type Culture Collection) was sequentially diluted for curve generation in aqueous solution containing equivalent amounts of DNA from uninfected mouse tissues. The following primers were used for *T. cruzi* genomic DNA, TCZ, GCTCTTGCCACACGGGTGC (forward), and CCAAGCAGCGGATAGTTCAGG (reverse); and for genomic *il-12p40*, GTAGAGGTGGACTGGACTCC (forward) and CAGATGTGAGTGGCTCAGAG (reverse).

Total RNA was isolated from spleens of WT and *phox* KO infected or non-infected mice and real-time RT-PCR was performed on an ABI PRISM 7900 sequence detection system (Applied Biosystems) using SYBR Green PCR Master Mix (Applied Biosystems) after RT of 1 µg RNA using SuperScript II reverse transcriptase (Invitrogen Life Technologies). The relative level of gene expression was determined by the comparative threshold cycle method as described by the manufacturer, whereby data for each sample were normalized to hypoxanthine phosphoribosyl transferase and expressed as a fold change compared with uninfected controls. The following primer pairs were used: for hypoxanthine phosphoribosyl transferase, GTTGGTTACAGGCCAGACTT-TGTTG (forward) and GAGGGTAGGCTGGCCTATAGGCT (reverse); *nos2*, CAGCTGGGCTGTACAAACCTT (forward) and CATTGGAAGTGAAGCGTTTCG (reverse).

Hepatic and pancreatic function

Serum AST and serum Amylase were measured in sera of infected and control animals using commercially available kits and following manufacturers instructions (KATAL, Belo Horizonte, MG, Brazil).

Determination of blood pressure by tail-cuff

After exposed for 5 minutes to a white lamp, WT and *phox* KO mice were placed in a plastic restrainer. Tail blood pressure (TBP) from the animals was measured using a pneumatic cuff placed in the base of the tail with a distally attached pulse sensor. Mice were allowed to adjust to this procedure three times a week for two weeks before experiments were performed. TBP values were recorded on a tail-cuff plethysmography Model MK-2000 using Windaq software to analyze the data. At least 10 good measurements for each animal were obtained per time point and the average of selected 5 best readings were used as TPB for an animal (n = 6 animals per group).

Determination of blood pressures by carotid catheterization

Mean arterial pressure (MAP) was recorded continuously in anesthetized animals by Biopac System (model MP150 A-CE, Biopac Systems, CA, USA) like described previously. In brief, mice were anesthetized by using urethane (1.2 g/kp) administered by intraperitoneal injection at different points after infection with *T. cruzi*. The adequacy of anesthesia was verified by the absence of a withdrawal response to nociceptive stimulation of a hindpaw. The left common carotid artery was exposed through a 1.0- to 1.5-cm midline incision in the ventral neck region. A catheter from polyethylene tubing (PE 5 Intramedic, Clay Adams, Becton Dickinson, Franklin Lakes, NJ, USA) was inserted approximately 0.25 cm into the common carotid artery and connected to pressure transducers. Supplemental doses of urethane (0.1 g/kg IV) were administered if necessary. The data were converted from

digital to numeric form using acquisition software. Data were processed by calculation of 10-min means of MAP variable. Results are expressed as means \pm SE. (measured in millimeters of mercury) of 2–6 animals per time point pooled from 3 independent experiments.

Treatment with iNOS inhibitors

Animals were treated with 1400SW, a NOS2 inhibitor (15 mg/Kg), i.p. on days 15 and 16 after infection with *T. cruzi*. On day 16, 1400W was administered 1 h before measuring MAP. During survival experiments, 1400W (20 mg/Kg) was administered i.p. daily divided in two doses or once a day beginning on day 13 after infection (a time found to not affect parasite control with NOS2 inhibition [35] and before MAP starts declining) for 10 days. Mice treated with vehicle were used as controls. Alternatively, animals were treated with aminoguanidine (1% w/v) in drinking water from day 13 after infection.

Statistics

The significance of differences between sample means was determined by Student's *t* test to compare WT to *phox* KO group or one-way ANOVA if *inf* KO animals were being compared as well. A mortality difference was tested using Mantel-Cox test and groups compared using one-way ANOVA. A value of $p < 0.05$ was considered significant.

Results

Mice deficient in functional NADPH oxidase control *T. cruzi* proliferation, but do not survive infection. *T. cruzi* infection is known to induce a strong oxidative stress in the host with high production of ROS and NO leading to nitration of serum and target organs proteins [24]. Results from our lab have shown that not only the ROS production is deficient in gp91^{phox} NADPH oxidase (*phox* KO) genetically deficient mice as described before [28], but the level of nitration of serum proteins induced by *T. cruzi* infection in *phox* KO mice is only 25% of that observed in WT controls (data not shown). To better investigate the role of ROS in *T. cruzi* infection in vivo, *phox* KO animals were inoculated with the Y strain of *T. cruzi*. Because this is a reticulotropic strain, it is more appropriate to evaluate the effects of ROS deficiency in phagocytes in vivo. WT and *phox* KO mice displayed similar parasitemia, which peaked around 9 days post-infection (Fig. 1A) and was subsequently controlled. In contrast, *ifn-γ* KO mice presented uncontrolled parasite counts throughout the infection. WT mice presented 60–70% of survival after day 50 of infection and all IFN- γ -deficient mice died by day 15 of infection. Surprisingly, *phox* KO animals exhibited high mortality when compared to WT controls, starting at day 15 and reaching 100% mortality by 21 days of infection (Fig. 1B). This unexpected result led us to investigate a possible parasite proliferation in tissues. Coherently with the parasitemia data, tissue parasitism was controlled by *phox* KO and WT groups at 15 days post-infection in spleens, livers, and heart; *ifn-γ* KO animals exhibited high parasite proliferation in these organs (Fig. 2).

phox KO and WT mice presented similar immune responses and pathology

The immune response from both WT and *phox* KO groups was analyzed. Both mouse strains displayed similar levels of TNF and IFN- γ in sera at 9 and 15 days post-infection (Fig. 3A). In addition, splenocytes from both groups produced expressive and equivalent levels of IFN- γ and TNF after 9 days of infection (Fig. 3B). Importantly, tissues from both animals exhibited similar quanti-




Figure 1. NADPH oxidase deficient-mice control parasitemia, but succumb to infection with *T. cruzi*. WT, *phox* KO and *ifn-γ* KO mice were infected with 1000 blood-born trypomastigotes of Y strain of *T. cruzi*. Parasitemia (A) and mortality (B) were accessed daily. (A) Points represent mean \pm SE of 5 animals per group of one from three different experiments performed with similar results. Asterisks represent $P < 0.05$ by Student's *t* test. (B) Mortality curve is pooled from three different experiments and $P < 0.05$ among all groups in the graph. doi:10.1371/journal.pntd.0001492.g001

tative and qualitative cellular infiltration in spleens, livers and hearts (not shown). Hepatic and pancreatic proofs were slightly increased after infection, but similar in both groups (Table 1).




Figure 2. *phox* KO mice control parasite proliferation in target organs. WT, *phox* KO and *ifn-γ* KO mice infected with *T. cruzi* were sacrificed on days 10 and 15 post-infection and tissue parasitism in spleen, heart and liver evaluated by real-time PCR as described in material and methods. Bars represent mean \pm SE of four animals per group. Arrows indicate $P < 0.05$ between WT and *phox* KO animals. The parasitism of *ifn-γ* KO group is statistically different from WT and *phox* groups in all organs and times analyzed, except for the heart at day 10 post-infection. doi:10.1371/journal.pntd.0001492.g002

NOx levels were exacerbated in *phox* KO mice with possible involvement in hemodynamic disturbances

Nitrate and nitrite (NOx) levels were evaluated in serum of infected mice. *Phox* KO mice exhibited about two fold higher levels when compared to WT-infected controls (Fig. 4A). Of note, NOx levels were increased in the *phox* KO group at the same time that mice began to die, about 15 days post-infection. The expression of *nos2* gene in the liver was measured by real-time RT-PCR and both WT and *phox* KO mice displayed similar levels of mRNA (Fig. 4B). Because NOx levels closely relate with pressoric regulation, the blood pressure was evaluated in the tail (TBP) using the non-invasive tail-cuff method and in the carotid artery by catheterization, at different time points (Fig. 5). When we evaluated the blood pressure in the tail, we observed that WT mice presented a good control of pressure variation as infection progressed, but *phox* KO mice exhibited dramatic oscillations of TBP after peak parasitemia (Fig. 5A). In order to have a more accurate picture of this phenomenon, we investigated the mean arterial pressure (MAP) in a central vessel, the carotid artery. As can be observed in figure 5B, the MAP of *phox* KO mice dropped from levels between 80–90 mmHg before infection to 70–60 mmHg by the time the NOx levels starts to increase in the serum, at day 8 post-infection, and further down as infection progressed. WT group displayed a good control of MAP till day 12 post-infection, but a drop in the blood pressure at day 14 to a level similar to that observed in the *phox* KO group occurred. While WT mice restored blood pressure to normal levels, *phox* KO counterparts were unable to restore physiological MAP (Fig. 5B).

In order to verify the role of NO produced by NOS2 in the drop of blood pressure and in mortality, *phox* KO mice were treated with 1400W, a selective inhibitor of NOS2. Injections with 1400W were able to inhibit NO levels in the blood (data not shown) and to restore blood pressure levels (Fig. 5C). However, animals treated daily (not shown) or every 12 hours with 1400W displayed similar mortality rates to that of control mice (Fig. 5D). We treated the animals with a less selective NOS2 inhibitor (aminoguanidine) in the drinking water (1% w/v) from day 13 of infection and no effect was observed on the mortality of *phox* KO infected mice (data not shown). These treatments did not impact the control of parasite proliferation in either WT or *phox* KO animals, nor changed the outcome of the disease in WT mice (data not shown).




Figure 3. WT and *phox* KO mice produce similar levels of IFN- γ and TNF. (A) WT and *phox* KO animals infected with *T. cruzi* were bled at days 10 and 15 post-infection for cytokine measurements. (B) Infected mice were sacrificed at 10 days post-infection and spleen cells isolated and cultured for 72 hours, when supernatants were harvested. IFN- γ and TNF were measured by ELISA as described in material and methods. Bars represent mean \pm SE of at least 4 animals per group. Experiment was repeated once with similar results. doi:10.1371/journal.pntd.0001492.g003

Discussion

The involvement of ROS in host resistance against infectious diseases is well known [36], especially for bacterial and fungal infections. However, while some reports suggest the involvement of ROS in protozoa infections [1,2,3,4,5], others fail to find a major effect of these radicals in control of infections with *L. major* [6], *T. gondii* [37] and *Plasmodium* [9]. Importantly, chronic granulomatous disease patients are known to suffer from severe bacterial and fungal infections [38], but rarely from severe protozoa infections [39]. Interestingly, data from our laboratory suggests that infection with *T. cruzi* can induce a strong oxidative state in the host with production of NO, ROS and superoxide causing nitration of proteins in serum and target tissue [24] (and data not shown). ROS is known to be produced by macrophages following *in vitro* *T. cruzi* infection and to be one of the major oxidative agents on *T. cruzi*, reducing its viability dramatically [19,21,40]. In this study, we investigated the role of ROS on *T. cruzi* infection *in vivo* and surprisingly we found an important physiological effect of ROS, unrelated to the control of parasite.

In the present study, we found that animals deficient in gp91^{phox} subunit of NADPH oxidase, a mouse model for chronic granulomatous disease [28], were able to efficiently control proliferation of Y strain of *T. cruzi*. Hence, parasitemia and parasite loads in spleen, liver and heart were similar in *phox* KO and WT mice. This result could suggest that ROS play a minor

role in restriction of protozoal infection during *in vivo* infections. On the other hand, when carefully examined *in vitro*, the effects of ROS on parasite control can be appreciated, especially the effect of peroxynitrite. For example, macrophage-derived ROS and peroxynitrite were found to cause major oxidative burden on *T. cruzi*, reducing its viability dramatically [19,21,40]. Indeed, the virulence of different parasite strains can be predicted by the expression of some enzymes involved in the parasite anti-oxidant network such as TcTS, TXN, TcMPX, TcAPX and FeSOD-A [25]. The fact that macrophage-derived ROS were found to have little involvement in parasite control in *phox* KO mice may be related to other mechanisms of resistance operating *in vivo* such as compensatory NO production, p47GTPases expression [17,18], CD8 T cells involvement [41] and alternative cellular sources of superoxide and peroxynitrite. Regarding this last point, we should indicate that normally, in activated macrophages, phagocyte-derived superoxide reacts with NO to yield peroxynitrite [21]; thus, in wild type animals superoxide from inflammatory cells plays a key role in NO-dependent cytotoxicity towards *T. cruzi* [20]. However, in the *phox* KO mice, the lack of macrophage-derived superoxide, increases the NO levels diffusing into the parasite, which in turn, inhibit the parasite mitochondrial respiration and secondarily enhance mitochondrial superoxide formation [25]. Overall, these processes lead to intramitochondrial formation of peroxynitrite and *T. cruzi* cytotoxicity. Indeed, the exceeding available NO in *phox* KO could be responsible for parasite control, including the formation of peroxynitrite in parasite mitochondria [20] or by NOX4, recently found in macrophages [42]. Higher levels of NO found in sera from *phox* KO mice could not be attributed to higher expression of NOS2. This could be explained simply by the fact that NO is not reacting with superoxide to yield peroxynitrite in *phox* KO. Another possibility is raised by the fact that superoxide facilitates uncoupling of NOS and oxidation of tetrahydrobiopterin, therefore in its absence NOS would be more active and produce more NO [43].

In addition to their anti-infection role, ROS are involved in enhancing TLR signaling. Recently, it was demonstrated that ROS production is activated by TLR signaling through MyD88 and via the p38 MAPKinase cascade [44]. After their production is activated by TLR-dependent or independent pathways, ROS are able to enhance TLR4 expression on the cell surface [45] and to strength NF- κ B activation [46]. The resistance to infection with *T. cruzi* is known to depend on appropriate MyD88 signaling [32] after stimulation of TLR2 and TLR9 [47], and TLR4 [48]. Although this function of

Table 1. Serum AST and amylase in WT and *phox* KO mice infected with *T. cruzi*.

		Days of infection:			
		0	8	12	15
AST ^a	WT	60.76 10.6	219.96 39.1	139.76 34.0	119.96 78.5
	<i>Phox</i> KO	73.06 5.0	207.66 35.9	225.56 22.5	136.36 87.0
Amylase ^b	WT	259.26 83.4	506.26 85.1	506.96 73.5	507.46 91.8
	<i>Phox</i> KO	246.46 113.3	520.66 161.9	526.16 47.9	486.66 146.2

Values from AST and amylase are combined from 3 independent experiments with n=3 for each independent experiment.

^aAST values are expressed in IU/L;

^bAmylase values are expressed in U/L.

doi:10.1371/journal.pntd.0001492.t001




Figure 4. Augmented NO_x levels in *phox*KO mice infected with *T. cruzi*, when compared to WT. (A) *T. cruzi*-infected mice were bled at 10 and 15 days post-infection and levels of nitrate and nitrite evaluated. Bars represent mean \pm SE of 4 animals per group. Asterisks indicate $P < 0.05$ by Student's *t* test. (B) Spleens from infected animals were harvested at 10 and 15 days post-infection and used for RNA extraction and real-time RT-PCR as described in material and methods. NOS2 expression was evaluated after normalization with HPRT constitutive gene. doi:10.1371/journal.pntd.0001492.g004

ROS could result in improved immunity to *T. cruzi*, it seemed to have no critical role in our system. Our results show that *phox* KO mice exhibited no immune impairment, producing equivalent amounts of IFN- γ and TNF in response to infection and presenting similar histopathology (data not shown) to their WT partners.

Surprisingly, despite the ability of *phox* KO mice to restrict *T. cruzi* infection and mount an efficient immune response, they

completely succumbed to infection by day 20 post-inoculation with *T. cruzi*. Further investigation showed that both WT and *phox* KO animals exhibited increased levels of NO_x in sera from day 8 to 15 post-infection. However, the levels of nitrogen intermediates were higher in *phox* KO at day 15, coinciding with the initiation of mortality. NO is produced by three different isoforms of nitric oxide synthase (NOS1 or neuronal NOS, NOS2 or inducible and




Figure 5. *T. cruzi*-infected *phox* KO mice display dramatic blood pressure variations. WT and *phox* KO animals were infected with *T. cruzi* and blood pressure evaluated in the tail (TBP) (A) or in the carotid artery (MAP) (B). Values represent mean \pm SE of 6 mice of one from two performed (A) or 2–6 mice per time point pooled from 3 independent experiments (B). Asterisks represent $P < 0.05$. (C) Drop in blood pressure is reverted by iNOS-specific inhibitor 1400SW. *phox* KO animals were infected with *T. cruzi* and blood pressure evaluated in the carotid artery (MAP) 16 days post-infection. Mice treated with 1400W received 15 mg/kg 24 and 1 hour before measurements. Values represent mean \pm SE of 4 mice of one from two performed. Asterisks represent $P < 0.05$. (D) 1400W did not revert mortality in *phox* KO mice. Mice were treated with 20 mg/kg of 1400W ip daily in single dose or divided in 2 doses starting on day 13 post-infection for 14 days. Mortality was accessed daily. Points represent mean cumulative mortality of 18–20 animals per group. Pool from 3 experiments performed with similar results (two experiments using one dose per day regime and one using two doses per day regime). doi:10.1371/journal.pntd.0001492.g005

NOS3 or endothelial) and is known to play a pleiotropic role in host physiology [49,50,51,52]. In addition to having potent antimicrobial properties, NO is involved in neurotransmission, gene expression and blood pressure regulation. For example, hyperproduction of NO has severe consequences to the host, being the cause of hypotension during septic shock [49]. During *T. cruzi* infection, uncontrolled immune response has been proven to be deleterious to the host, as is the case of infection in the absence of IL10 [53,54]. In *phox* KO animals excess NO was associated with peripheral blood pressure variations, not observed in WT controls and, more importantly, with early and permanent drop in central MAP. An important drop in MAP of WT animals was also observed, but this hypotension happened later than in *phox* KO mice and was transitory, lasting for no longer than 3 days. Although WT mice showed death rate of 40% starting on day 20 post-infection, we do not think this mortality is associated with the levels of NO since it starts after full recovery of blood pressure levels. From this study, we can conclude also that several factors might be involved in death associated with experimental *T. cruzi* infection. For example, the fact that treatment with 1400W prevented blood pressure drop in *phox* KO mice implicates NOS2. However, treatment was not able to prevent death. Inhibition of NOS using NOS inhibitors early in *T. cruzi* infection results in higher mortality due to infection [24,55]. In contrast, treatment of *T. cruzi*-infected mice with NOS inhibitors in the chronic phase of the infection (Tulahuen strain) was not detrimental to the host's ability to control parasitism [35]. In addition, NOS2-deficient animals, in contrast to *ifn-γ* KO mice, can survive if treated with suboptimal doses of benzonidazole during peak of parasitemia even if the drug is withdrawn after parasite control [14]. We followed parasitemia in animals treated with NOS2 inhibitors after parasitemia was controlled and we did not observe recrudescence of parasite proliferation. These data suggest that NO may have an important role especially in the acute phase of the infection, in contrast to chronic phase when other IFN- γ -dependent mechanism controls the infection. The fact that NOS2 inhibition, although improving blood pressure, did not prevent mortality in our experiment could suggest that the cause of death may be multi-factorial possibly involving changes in hematological parameters (infection associated anemia and leucopenia) [56] or cardiac function [57,58] and demands further investigation. However, data from shock models show that restoring blood pressures to normal levels may not rescue animals from death. The

reason for this failure would be that the iNOS inhibition enhances the accumulation of activated leukocytes into vital organs, thus increasing tissue lesions. Also, inhibition of iNOS reduces the perfusion of the organs [59,60,61,62,63].

Another very interesting side of ROS actions started to be depicted recently. ROS have been shown to regulate vasoactive properties of NO. Nitric oxide is known to react with the heme group of guanylate cyclase activating the production of cGMP that promotes vasodilation [64]. Accordingly, some inhibitors of guanylate cyclase, such as methylene blue, induce ROS production. In addition, ROS derived from endothelial NADPH oxidase containing gp91^{phox} is a potent vasoconstrictor because it scavenges NO before NO activates guanylate cyclase [29,30]. Hence, one unifying hypothesis to explain an important part of our observations is that *T. cruzi* infection stimulates a strong production of NO and *phox* KO animals cannot produce ROS in order to counteract the systemic effect of NO. Coherently, despite elevated levels of NOx detected in sera of *phox* KO animals, they expressed similar levels of NOS2 by real-time RT-PCR in the spleen. This fact and the finding that IFN- γ and TNF are not increased in *phox* KO animals suggest that it is not likely that elevated NOx in serum is due to augmented production, but may be related to impaired ROS production and its role in scavenging NO. We propose that the reaction of ROS and NO to generate peroxynitrite, in addition to strengthening the killing effects of NO by augmenting its oxidative properties [21], has an important role in regulating NO signaling and its systemic effects during *T. cruzi* infection.

Acknowledgments

We are indebted to Antonio Mesquita Vaz for excellent animal care, to Dr. Tom Nuttman for providing 1400W, to Dr. Yara Quercia Vieira and Mr. Edgar Sleifer for reagents and to Dr. Fabiana Romão Machado for help in experiments and useful suggestions.

Author Contributions

Conceived and designed the experiments: HCS CZGL FQC MMT RR LQV. Performed the experiments: HCS CZGL JPM LU WLT. Analyzed the data: HCS CZGL FQC MMT RR LQV. Contributed reagents/materials/analysis tools: MJCS ROA JCFAF AJR FQC MMT. Wrote the paper: HCS CZGL RR LQV.

References

- Murray HW, Rubin BY, Carriero SM, Harris AM, Jaffee EA (1985) Human mononuclear phagocyte antiparasitic mechanisms: oxygen-dependent vs oxygen-independent activity against intracellular *Toxoplasma gondii*. *J Immunol* 134: 1982–1988.
- Shrestha SP, Tomita T, Weiss LM, Orloffsky A (2006) Proliferation of *Toxoplasma gondii* in inflammatory macrophages in vivo is associated with diminished oxygen radical production in the host cell. *Int J Parasitol* 36: 433–441.
- Golenser J, Kamyli M, Tsafack A, Marva E, Cohen A, et al. (1992) Correlation between destruction of malarial parasites by polymorphonuclear leukocytes and oxidative stress. *Free Radic Res Commun* 17: 249–262.
- Blos M, Schleicher U, Soares Rocha FJ, Meissner U, Rollinghoff M, et al. (2003) Organ-specific and stage-dependent control of *Leishmania major* infection by inducible nitric oxide synthase and phagocyte NADPH oxidase. *Eur J Immunol* 33: 1224–1234.
- Murray HW, Nathan CF (1999) Macrophage microbicidal mechanisms in vivo: reactive nitrogen versus oxygen intermediates in the killing of intracellular visceral *Leishmania donovani*. *J Exp Med* 189: 741–746.
- Assreuy J, Cunha FQ, Eppertlein M, Noronha-Dutra A, O'Donnell CA, et al. (1994) Production of nitric oxide and superoxide by activated macrophages and killing of *Leishmania major*. *Eur J Immunol* 24: 672–676.
- Ding M, Kwok LY, Schluter D, Clayton C, Soldati D (2004) The antioxidant systems in *Toxoplasma gondii* and the role of cytosolic catalase in defence against oxidative injury. *Mol Microbiol* 51: 47–61.
- Pewes KA, Barr SD, Gedamu L (2003) Iron superoxide dismutases targeted to the glycosomes of *Leishmania chagasi* are important for survival. *Infect Immun* 71: 5910–5920.
- Sobolewski P, Gramaglia I, Frangos JA, Intaglietta M, van der Heyde H (2005) *Plasmodium berghei* resists killing by reactive oxygen species. *Infect Immun* 73: 6704–6710.
- Brunet LR (2001) Nitric oxide in parasitic infections. *Int Immunopharmacol* 1: 1457–1467.
- Green SJ, Scheller LF, Marletta MA, Seguin MC, Klotz FW, et al. (1994) Nitric oxide: cytokine-regulation of nitric oxide in host resistance to intracellular pathogens. *Immunol Lett* 43: 87–94.
- Adams LB, Dinauer MC, Morgenstern DE, Krahenbuhl JL (1997) Comparison of the roles of reactive oxygen and nitrogen intermediates in the host response to *Mycobacterium tuberculosis* using transgenic mice. *Tuber Lung Dis* 78: 237–246.
- Cummings KL, Tarleton RL (2004) Inducible nitric oxide synthase is not essential for control of *Trypanosoma cruzi* infection in mice. *Infect Immun* 72: 4081–4089.
- Michailowsky V, Silva NM, Rocha CD, Vieira LQ, Lannes-Vieira J, et al. (2001) Pivotal role of interleukin-12 and interferon-gamma axis in controlling tissue parasitism and inflammation in the heart and central nervous system during *Trypanosoma cruzi* infection. *Am J Pathol* 159: 1723–1733.
- Scharton-Kersten TM, Yap G, Magram J, Sher A (1997) Inducible nitric oxide is essential for host control of persistent but not acute infection with the intracellular pathogen *Toxoplasma gondii*. *J Exp Med* 185: 1261–1273.

16. Huang H, Chan J, Wittner M, Jelicks LA, Morris SA, et al. (1999) Expression of cardiac cytokines and inducible form of nitric oxide synthase (NOS2) in *Trypanosoma cruzi*-infected mice. *J Mol Cell Cardiol* 31: 75–88.
17. Koga R, Hamano S, Kuwata H, Aarashi K, Ogawa M, et al. (2006) TLR-dependent induction of IFN- β mediates host defense against *Trypanosoma cruzi*. *J Immunol* 177: 7059–7066.
18. Santiago HC, Feng CG, Bafica A, Roffe E, Arantes RM, et al. (2005) Mice deficient in LRG-47 display enhanced susceptibility to *Trypanosoma cruzi* infection associated with defective hemopoiesis and intracellular control of parasite growth. *J Immunol* 175: 8165–8172.
19. Alvarez MN, Piacenza L, Irigoin F, Peluffo G, Radi R (2004) Macrophage-derived peroxynitrite diffusion and toxicity to *Trypanosoma cruzi*. *Arch Biochem Biophys* 432: 222–232.
20. Piacenza L, Alvarez MN, Peluffo G, Radi R (2009) Fighting the oxidative assault: the *Trypanosoma cruzi* journey to infection. *Curr Opin Microbiol* 12: 415–421.
21. Alvarez MN, Peluffo G, Piacenza L, Radi R (2010) Intraphagosomal peroxynitrite as a macrophage-derived cytotoxin against internalized *Trypanosoma cruzi*: Consequences for oxidative killing and role of microbial peroxidases in infectivity. *J Biol Chem* 285: 6627–6640.
22. Gazzinelli RT, Oswald IP, Hieny S, James SL, Sher A (1992) The microbicidal activity of interferon- γ -treated macrophages against *Trypanosoma cruzi* involves an L-arginine-dependent, nitrogen oxide-mediated mechanism inhibited by interleukin-10 and transforming growth factor- β . *Eur J Immunol* 22: 2501–2506.
23. Dhiman M, Estrada-Franco JG, Pando JM, Ramirez-Aguilar FJ, Spratt H, et al. (2009) Increased myeloperoxidase activity and protein nitration are indicators of inflammation in patients with Chagas' disease. *Clin Vaccine Immunol* 16: 660–666.
24. Naviliat M, Gualco G, Cayota A, Radi R (2005) Protein 3-nitrotyrosine formation during *Trypanosoma cruzi* infection in mice. *Braz J Med Biol Res* 38: 1825–1834.
25. Piacenza L, Zago MP, Peluffo G, Alvarez MN, Basombrio MA, et al. (2009) Enzymes of the antioxidant network as novel determiners of *Trypanosoma cruzi* virulence. *Int J Parasitol* 39: 1455–1464.
26. Piacenza L, Peluffo G, Alvarez MN, Kelly JM, Wilkinson SR, et al. (2008) Peroxiredoxins play a major role in protecting *Trypanosoma cruzi* against macrophage- and endogenously-derived peroxynitrite. *Biochem J* 410: 359–368.
27. Radi R (2004) Nitric oxide, oxidants, and protein tyrosine nitration. *Proc Natl Acad Sci U S A* 101: 4003–4008.
28. Pollock JD, Williams DA, Gifford MA, Li LL, Du X, et al. (1995) Mouse model of X-linked chronic granulomatous disease, an inherited defect in phagocyte superoxide production. *Nat Genet* 9: 202–209.
29. Gorlach A, Brandes RP, Nguyen K, Amidi M, Dehghani F, et al. (2000) A gp91phox containing NADPH oxidase selectively expressed in endothelial cells is a major source of oxygen radical generation in the arterial wall. *Circ Res* 87: 26–32.
30. Jung O, Schreiber JG, Geiger H, Pedrazzini T, Busse R, et al. (2004) gp91phox-containing NADPH oxidase mediates endothelial dysfunction in renovascular hypertension. *Circulation* 109: 1795–1801.
31. Dalton DK, Pitts-Mekie S, Keshav S, Figari JS, Bradley A, et al. (1993) Multiple defects of immune cell function in mice with disrupted interferon- γ genes. *Science* 259: 1739–1742.
32. Campos MA, Closel M, Valente EP, Cardoso JE, Akira S, et al. (2004) Impaired production of proinflammatory cytokines and host resistance to acute infection with *Trypanosoma cruzi* in mice lacking functional myeloid differentiation factor 88. *J Immunol* 172: 1711–1718.
33. Ding AH, Nathan CF, Stuehr DJ (1988) Release of reactive nitrogen intermediates and reactive oxygen intermediates from mouse peritoneal macrophages. Comparison of activating cytokines and evidence for independent production. *J Immunol* 141: 2407–2412.
34. Cummings KL, Tarleton RL (2003) Rapid quantitation of *Trypanosoma cruzi* in host tissue by real-time PCR. *Mol Biochem Parasitol* 129: 53–59.
35. Saefel M, Fleischer B, Hoerauf A (2001) Stage-dependent role of nitric oxide in control of *Trypanosoma cruzi* infection. *Infect Immun* 69: 2252–2259.
36. Nathan C, Shiloh MU (2000) Reactive oxygen and nitrogen intermediates in the relationship between mammalian hosts and microbial pathogens. *Proc Natl Acad Sci U S A* 97: 8841–8848.
37. Alexander J, Scharton-Kersten TM, Yap G, Roberts CW, Liew FY, et al. (1997) Mechanisms of innate resistance to *Toxoplasma gondii* infection. *Philos Trans R Soc Lond B Biol Sci* 352: 1355–1359.
38. Heyworth PG, Cross AR, Curmutte JT (2003) Chronic granulomatous disease. *Curr Opin Immunol* 15: 578–584.
39. Asensi V, Tricas L, Meana A, Roos D, Carton JA, et al. (2000) Visceral leishmaniasis and other severe infections in an adult patient with p47-phox-deficient chronic granulomatous disease. *Infection* 28: 171–174.
40. Piacenza L, Irigoin F, Alvarez MN, Peluffo G, Taylor MC, et al. (2007) Mitochondrial superoxide radicals mediate programmed cell death in *Trypanosoma cruzi*: cytoprotective action of mitochondrial iron superoxide dismutase overexpression. *Biochem J* 403: 323–334.
41. Padilla AM, Bustamante JM, Tarleton RL (2009) CD8 $^{+}$ T cells in *Trypanosoma cruzi* infection. *Curr Opin Immunol* 21: 385–390.
42. Lee CF, Qiao M, Schroder K, Zhao Q, Asmis R (2010) Nox4 is a novel inducible source of reactive oxygen species in monocytes and macrophages and mediates oxidized low density lipoprotein-induced macrophage death. *Circ Res* 106: 1489–1497.
43. Sun J, Druhan LJ, Zweier JL (2010) Reactive oxygen and nitrogen species regulate inducible nitric oxide synthase function shifting the balance of nitric oxide and superoxide production. *Arch Biochem Biophys* 494: 130–137.
44. Laroux FS, Romero X, Wetzler L, Engel P, Terhorst C (2005) Cutting edge: MyD88 controls phagocyte NADPH oxidase function and killing of gram-negative bacteria. *J Immunol* 175: 5596–5600.
45. Powers KA, Szasz K, Khadaroo RG, Tawadros PS, Marshall JC, et al. (2006) Oxidative stress generated by hemorrhagic shock recruits Toll-like receptor 4 to the plasma membrane in macrophages. *J Exp Med* 203: 1951–1961.
46. Ryan KA, Smith MF, Jr., Sanders MK, Ernst PB (2004) Reactive oxygen and nitrogen species differentially regulate Toll-like receptor 4-mediated activation of NF- κ B and interleukin-8 expression. *Infect Immun* 72: 2123–2130.
47. Bafica A, Santiago HC, Goldszmid R, Ropert C, Gazzinelli RT, et al. (2006) Cutting edge: TLR9 and TLR2 signaling together account for MyD88-dependent control of parasitemia in *Trypanosoma cruzi* infection. *J Immunol* 177: 3515–3519.
48. Oliveira AC, Peixoto JR, de Arruda LB, Campos MA, Gazzinelli RT, et al. (2004) Expression of functional TLR4 confers proinflammatory responsiveness to *Trypanosoma cruzi* glycoinositolphospholipids and higher resistance to infection with *T. cruzi*. *J Immunol* 173: 5688–5696.
49. Assreuy J (2006) Nitric oxide and cardiovascular dysfunction in sepsis. *Endocr Metab Immune Disord Drug Targets* 6: 165–173.
50. Beck KF, Eberhardt W, Frank S, Huwiler A, Messmer UK, et al. (1999) Inducible NO synthase: role in cellular signalling. *J Exp Biol* 202: 645–653.
51. Bogdan C (2001) Nitric oxide and the immune response. *Nat Immunol* 2: 907–916.
52. MacMicking J, Xie QW, Nathan C (1997) Nitric oxide and macrophage function. *Annu Rev Immunol* 15: 323–350.
53. Holscher C, Mohrs M, Dai WJ, Kohler G, Ryffel B, et al. (2000) Tumor necrosis factor α -mediated toxic shock in *Trypanosoma cruzi*-infected interleukin 10-deficient mice. *Infect Immun* 68: 4075–4083.
54. Hunter CA, Ellis-Neyes LA, Slifer T, Kanaly S, Grunig G, et al. (1997) IL-10 is required to prevent immune hyperactivity during infection with *Trypanosoma cruzi*. *J Immunol* 158: 3311–3316.
55. Vespa GN, Cunha FQ, Silva JS (1994) Nitric oxide is involved in control of *Trypanosoma cruzi*-induced parasitemia and directly kills the parasite in vitro. *Infect Immun* 62: 5177–5182.
56. Malvezi AD, Cecchini R, de Souza F, Tadokoro CE, Rizzo LV, et al. (2004) Involvement of nitric oxide (NO) and TNF- α in the oxidative stress associated with anemia in experimental *Trypanosoma cruzi* infection. *FEMS Immunol Med Microbiol* 41: 69–77.
57. Durand JL, Mukherjee S, Commodari F, De Souza AP, Zhao D, et al. (2009) Role of NO synthase in the development of *Trypanosoma cruzi*-induced cardiomyopathy in mice. *Am J Trop Med Hyg* 80: 782–787.
58. Durand JL, Tang B, Gutstein DE, Petkova S, Teixeira MM, et al. (2006) Dyskinesia in Chagasic myocardium: centerline analysis of wall motion using cardiac-gated magnetic resonance images of mice. *Magn Reson Imaging* 24: 1051–1057.
59. Avontuur JA, Bruining HA, Ince C (1995) Inhibition of nitric oxide synthesis causes myocardial ischemia in endotoxemic rats. *Circ Res* 76: 418–425.
60. Henderson JL, Statman R, Cunningham JN, Cheng W, Damiani P, et al. (1994) The effects of nitric oxide inhibition on regional hemodynamics during hyperdynamic endotoxemia. *Arch Surg* 129: 1271–1274. discussion 1275.
61. Hickey MJ, Sharkey KA, Sihota EG, Reinhardt PH, MacMicking JD, et al. (1997) Inducible nitric oxide synthase-deficient mice have enhanced leukocyte-endothelium interactions in endotoxemia. *FASEB J* 11: 955–964.
62. Laubach VE, Shesely EG, Smithies O, Sherman PA (1995) Mice lacking inducible nitric oxide synthase are not resistant to lipopolysaccharide-induced death. *Proc Natl Acad Sci U S A* 92: 10688–10692.
63. MacMicking JD, Nathan C, Hom G, Chartrain N, Fletcher DS, et al. (1995) Altered responses to bacterial infection and endotoxic shock in mice lacking inducible nitric oxide synthase. *Cell* 81: 641–650.
64. Ignarro LJ (1990) Nitric oxide. A novel signal transduction mechanism for transcellular communication. *Hypertension* 16: 477–483.